

Notes for *Analyse 1* and *Analyse 2*

Horia Cornean, 30/04/2012.

Contents

1	The natural topology of a metric space	2
2	Compact and sequentially compact sets	3
2.1	Compact implies sequentially compact	4
2.2	Sequentially compact implies compact	4
2.3	The Heine-Borel Theorem	6
3	Continuous functions on metric spaces	7
4	Banach's fixed point theorem	9
5	Local existence and uniqueness for first order ODE's	9
5.1	Spaces of bounded/continuous functions	10
5.2	The main theorem	11
6	The inverse function theorem	13
7	The implicit function theorem	16
8	Brouwer's fixed point theorem	18
9	Schauder's fixed point theorem	22
10	Kakutani's fixed point theorem	23
11	Existence of Nash equilibrium for finite games with two players	24
12	The Hairy Ball Theorem	25
13	The Jordan Curve Theorem	27
13.1	Some preparatory results	27
13.2	The main theorem	31

These notes are strongly inspired by the books *Principles of Mathematical Analysis* by Walter Rudin and *Topology from the Differentiable Viewpoint* by John Milnor. Some of the theorems below can be formulated in a more general setting than the one of metric spaces, but the metric space structure brings important simplifications and clarity. Fundamental results like the Brouwer, Schauder and Kakutani Fixed Point Theorems, the Hairy Ball Theorem, the Tietze Extension Theorem, and the Jordan Curve Theorem are not in the curriculum, but this does not make them less important. All proofs are quite detailed and self-contained, and are at the level of hard-working second year undergraduate students.

The first two chapters deal with point set topology in metric spaces. In particular we prove the equivalence between compact and sequentially compact sets in metric spaces, and the Heine-Borel Theorem in Euclidian spaces. We assume the Bolzano-Weierstrass Theorem to be known.

Chapter three deals with continuous functions on metric spaces. We show the equivalence between continuity, sequential continuity, and 'returning open sets into open sets'. We show that a continuous function defined on a compact set is uniformly continuous.

Chapter four proves the Banach Fixed Point Theorem. Chapter five is based on the previous one and investigates the local existence and uniqueness of solutions to first order differential equations.

Chapter six contains the Inverse Function Theorem. Its proof is based on Banach's Fixed Point Theorem. Chapter seven deals with the Implicit Function Theorem, whose proof is shown to be a consequence of the Inverse Function Theorem.

Chapter eight contains the proof of the Brouwer Fixed Point Theorem. We follow the strategy of C.A. Rogers from the paper *A Less Strange Version of Milnor's Proof of Brouwer's Fixed Point Theorem*, appeared in Amer. Math. Monthly. **87** 525-527 (1980). We give many more details and the presentation is completely analytic and self-contained, based on the previous six chapters. We also prove that any convex body is homeomorphic with the closed unit ball.

Chapter nine contains the Schauder Fixed Point Theorem, presented as a consequence of Brouwer's Fixed Point Theorem.

Chapter ten presents the Kakutani Fixed Point Theorem. Its proof is an adaptation of that of S. Kakutani in *A generalization of Brouwer's fixed point theorem*, Duke Mathematical Journal **8**(3), 457-459 (1941). This theorem is another consequence of Brouwer's Fixed Point Theorem.

Chapter eleven contains the proof of the existence of a Nash equilibrium for a finite game with two players, based on Kakutani's theorem. The original paper of J. Nash entitled *Non-cooperative games*, Annals of Math. **54** (2), 286-295 (1951) had a slightly different proof, based on Brouwer's fixed point theorem.

Chapter twelve contains an analytic proof of the Hairy Ball Theorem, and it follows the strategy used by J. Milnor in the paper *Analytic proofs of the hairy ball theorem and the Brouwer fixed point theorem*, appeared in Amer. Math. Monthly **85**, 521-524 (1978).

Chapter thirteen contains the Jordan Curve Theorem and is inspired by a paper of R. Maehara entitled *The Jordan curve theorem via the Brouwer fixed point theorem*, which appeared in Amer. Math. Monthly **91**(10), 641-643 (1984). We give many more details and in particular, we prove a simple version of the Tietze Extension Theorem in \mathbb{R}^2 based on an extension formula due to Hausdorff.

1 The natural topology of a metric space

Let (X, d) be a metric space. We define the open ball of radius $r > 0$ and center at $a \in X$ the set $B_r(a) := \{x \in X : d(x, a) < r\}$.

Given a set $A \subset X$ and $a \in A$, we say that a is an interior point of A if there exists $r > 0$ such that $B_r(a) \subset A$. The set of all interior points of A is denoted by $\text{Int}(A)$. We say that A is an open set if all its points are interior points, i.e. $\text{Int}(A) = A$. By convention, the empty set \emptyset is open.

Lemma 1.1. *Any ball $B_r(a)$ is an open set.*

Proof. Let $x_0 \in B_r(a)$. We have that $d(x_0, a) < r$. Define $r_0 := (r - d(x_0, a))/2 > 0$. Then for all $x \in B_{r_0}(x_0)$ we have that $d(x, x_0) < r_0$ and:

$$d(x, a) \leq d(x, x_0) + d(x_0, a) < (r - d(x_0, a))/2 + d(x_0, a) = (r + d(x_0, a))/2 < r,$$

which shows that $B_{r_0}(x_0) \subset B_r(a)$. Thus $B_r(a)$ has only interior points. \square

Lemma 1.2.

- (i). Let $\{V_\alpha\}_{\alpha \in \mathcal{F}}$ be an arbitrary collection of open sets. Then $A := \cup_{\alpha} V_\alpha$ is open.
- (ii). Let $\{V_j\}_{j=1}^n$ be a finite collection of open sets. Then $B := \cap_{j=1}^n V_j$ is open.

Proof. We start with (i). Let $a \in \cup_{\alpha} V_\alpha$. There must exist $\alpha_a \in \mathcal{F}$ such that $a \in V_{\alpha_a}$. Since V_{α_a} is open, there exists $r_a > 0$ such that

$$B_{r_a}(a) \subset V_{\alpha_a} \subset \cup_{\alpha} V_\alpha = A$$

hence a is an interior point of A .

We continue with (ii). Let $a \in \cap_{j=1}^n V_j$. Thus $a \in V_j$ for all j . Hence there exists $r_j > 0$ such that $B_{r_j}(a) \subset V_j$. Let $r := \min\{r_1, \dots, r_n\} > 0$. Thus $B_r(a) \subset B_{r_j}(a) \subset V_j$ for all j , hence $B_r(a) \subset B$ and we are done. \square

We say that a set $A \subset X$ is closed if $A^c := \{x \in X : x \notin A\}$ is open. Given a set $B \subset X$ and $b \in X$, we say that b is an adherent point of B if there exists a sequence $\{x_n\}_{n \geq 1} \subset B$ such that $x_n \in B_{\frac{1}{n}}(b)$ and $\lim_{n \rightarrow \infty} x_n = b$. The set of all adherent points of B is denoted by \overline{B} .

Theorem 1.3. Let $B \subset X$. Then $B \subset \overline{B}$. Moreover, $B = \overline{B}$ if and only if B is closed.

Proof. If $a \in B$ we can define the constant sequence $x_n = a \in B$ which converges to a , thus $a \in \overline{B}$ and $B \subset \overline{B}$.

Now assume that $B = \overline{B}$. We want to prove that B is closed, i.e. B^c is open. Let $a \in B^c = \overline{B}^c$. Then a is not an adherent point, which means that there exists $\epsilon > 0$ such that no point of B lies in the ball $B_\epsilon(a)$. In other words, $B_\epsilon(a) \subset B^c$, hence B^c is open.

Now assume that B is closed. We want to prove that $B = \overline{B}$. Assume that this is not true; it would imply the existence of a point $b \in \overline{B}$ such that $b \in B^c$. Since B^c is open, there exists $\epsilon > 0$ such that $B_\epsilon(b) \subset B^c$, i.e. $B_\epsilon(b) \cap B = \emptyset$. But this is incompatible with $b \in \overline{B}$. \square

2 Compact and sequentially compact sets

Definition 2.1. Let A be a subset of a metric space (X, d) . Let \mathcal{F} be an arbitrary set of indices, and consider the family of sets $\{\mathcal{O}_\alpha\}_{\alpha \in \mathcal{F}}$, where each $\mathcal{O}_\alpha \subseteq X$ is open. This family is called an open covering of A if $A \subseteq \cup_{\alpha \in \mathcal{F}} \mathcal{O}_\alpha$.

Definition 2.2. Assume that $\{\mathcal{O}_\alpha\}_{\alpha \in \mathcal{F}}$ is an open covering of A . If \mathcal{F}' is a subset of \mathcal{F} , we say that $\{\mathcal{O}_\alpha\}_{\alpha \in \mathcal{F}'}$ is a subcovering if we still have the property $A \subseteq \cup_{\alpha \in \mathcal{F}'} \mathcal{O}_\alpha$. A subcovering is called finite, if \mathcal{F}' contains finitely many elements.

Definition 2.3. Let A be a subset of a metric space (X, d) . Then we say that A is covered by a finite ϵ -net if there exists a natural number $N_\epsilon < \infty$ and the points $\{\mathbf{x}_1, \dots, \mathbf{x}_{N_\epsilon}\} \subseteq A$ such that $A \subseteq \cup_{j=1}^{N_\epsilon} B_\epsilon(\mathbf{x}_j)$.

Definition 2.4. A subset $A \subset X$ is called compact, if from any open covering of A one can extract a finite subcovering.

Definition 2.5. $A \subset X$ is called sequentially compact if from any sequence $\{x_n\}_{n \geq 1} \subseteq A$ one can extract a subsequence $\{x_{n_k}\}_{k \geq 1}$ which converges to some point $x_\infty \in A$.

We will see that in metric spaces the two notions of compactness are equivalent.

2.1 Compact implies sequentially compact

We begin with two lemmas:

Lemma 2.6. Assume that the sequence $\{x_n\}_{n \geq 1} \subset A$ has a range consisting of finitely many points. Then it admits a convergent subsequence whose limit is one of the elements in the range.

Proof. Assume that the range of the sequence consists of the distinct points a_1, a_2, \dots, a_N . At least one of these points, say a_1 , is taken infinitely many times by the sequence elements. Denote by n_k (with $k \geq 1$) the increasing sequence of indices for which $x_{n_k} = a_1$. This defines our convergent subsequence. \square

Lemma 2.7. Assume that the sequence $\{x_n\}_{n \geq 1} \subset A$ has an accumulation point $a \in A$, i.e. for every $\epsilon > 0$ there exists some $x_{n_\epsilon} \neq a$ such that $x_{n_\epsilon} \in B_\epsilon(a)$. Then $\{x_n\}_{n \geq 1}$ admits a convergent subsequence whose limit is a .

Proof. Since a is an accumulation point, then for every $k \geq 1$ there exists some element x_{n_k} such that $0 < d(x_{n_k}, a) < 1/k$. This shows that $\{x_{n_k}\}_{k \geq 1}$ converges to a . \square

Theorem 2.8. Let $A \subseteq X$ be compact. Then A is sequentially compact.

Proof. We will assume the opposite, i.e. there exists a sequence $\{x_n\}_{n \geq 1}$ with no convergent subsequence in A . Such a sequence must have an infinite number of distinct points in the range, due to Lemma 2.6. Moreover, we can assume that $\{x_n\}_{n \geq 1}$ has no accumulation points in A (otherwise such a point would be the limit of a subsequence according to Lemma 2.7).

Since no $x \in A$ can be an accumulation point for $\{x_n\}_{n \geq 1}$, there exists $\epsilon_x > 0$ such that the ball $B_{\epsilon_x}(x)$ contains at most one element of the range of $\{x_n\}_{n \geq 1}$.

Clearly, $\{B_{\epsilon_x}(x)\}_{x \in A}$ is an open covering for A . Because A is compact, we can extract a finite subcovering from it:

$$A \subseteq \bigcup_{j=1}^N B_{\epsilon_{x_j}}(x_j), \quad N < \infty.$$

Now remember that $\{x_n\}_{n \geq 1} \subseteq A \subseteq \bigcup_{j=1}^N B_{\epsilon_{x_j}}(x_j)$ and at the same time, there are at most N distinct points of this sequence in the union $\bigcup_{j=1}^N B_{\epsilon_{x_j}}(x_j)$. We conclude that $\{x_n\}_{n \geq 1}$ can only have a finite number of distinct points in its range, thus it must admit a convergent subsequence according to Lemma 2.6. This contradicts our hypothesis. \square

2.2 Sequentially compact implies compact

The proof of this fact is slightly more complicated. We need two preparatory results:

Proposition 2.9. Let A be a sequentially compact set. Then for every $\epsilon > 0$, A can be covered by a finite ϵ -net (see Definition 2.3).

Proof. If A contains finitely many points, then the proof is obvious, thus we may assume that $\#(A) = \infty$.

Now suppose that there exists some $\epsilon_0 > 0$ such that A cannot be covered by a finite ϵ_0 -net. This means that for any N points of A , $\{x_1, \dots, x_N\}$, we have:

$$A \not\subseteq \bigcup_{j=1}^N B_{\epsilon_0}(x_j). \quad (2.1)$$

We will now construct a sequence with elements in A which cannot have a convergent subsequence. Choose an arbitrary point $x_1 \in A$. We know from (2.1), for $N = 1$, that we can find $x_2 \in A$ such that $x_2 \in A \setminus B_{\epsilon_0}(x_1)$. This means that $d(x_1, x_2) \geq \epsilon_0$. We use (2.1) again, for $N = 2$, in order to get a point $x_3 \in A \setminus [B_{\epsilon_0}(x_1) \cup B_{\epsilon_0}(x_2)]$. This means that $d(x_3, x_1) \geq \epsilon_0$ and $d(x_3, x_2) \geq \epsilon_0$. Thus we can continue with this procedure and construct a sequence $\{x_n\}_{n \geq 1} \subseteq A$ which obeys

$$d(x_j, x_k) \geq \epsilon_0, \quad j \neq k.$$

In other words, we constructed a sequence in A which cannot have a Cauchy subsequence. This contradicts Definition 2.5. \square

The second result states that a compact set is bounded:

Lemma 2.10. *Let A be a (sequentially) compact set. Then there exists a ball which contains A .*

Proof. We know that A can be covered by any finite ϵ -net; choose $\epsilon = 1$. Then there exist N points of A denoted by $\{x_1, \dots, x_N\}$ such that $A \subseteq \bigcup_{j=1}^N B_1(x_j)$.

Denote by $R = \max\{1 + d(x_j, x_k) : 1 \leq j, k \leq N\}$. Then we have $B_1(x_j) \subseteq B_R(x_1)$ for every j , thus $A \subseteq B_R(x_1)$ and we are done. \square

Let us now prove the theorem:

Theorem 2.11. *Assume that $A \subseteq X$ is sequentially compact. Then A is compact.*

Proof. Consider an arbitrary open covering of A :

$$A \subseteq \bigcup_{\alpha \in \mathcal{F}} \mathcal{O}_\alpha.$$

We will show that we can extract a finite subcovering from it.

For every $x \in A$, there exists at least one open set $\mathcal{O}_{\alpha(x)}$ such that $x \in \mathcal{O}_{\alpha(x)}$. Because $\mathcal{O}_{\alpha(x)}$ is open, we can find $\epsilon > 0$ such that $B_\epsilon(x) \subseteq \mathcal{O}_{\alpha(x)}$.

For a fixed x , we define the set

$$E_x := \{r > 0 : \text{there exists } \alpha \in \mathcal{F} \text{ such that } B_r(x) \subseteq \mathcal{O}_\alpha\} \subset \mathbb{R}.$$

From the above argument we conclude that no E_x is empty. Moreover, if $r \in E_x$, then the open interval $(0, r)$ is included in E_x .

If for some x in A we have an unbounded E_x , it follows that for every $r > 0$ we can find some open set \mathcal{O}_α such that $B_r(x) \subseteq \mathcal{O}_\alpha$. But if r is chosen to be large enough, it will contain the ball we constructed in Lemma 2.10, thus \mathcal{O}_α will also contain A . In this case we found our subcovering, which consists of just one open set.

It follows that we may assume that all the sets E_x are bounded intervals admitting a positive and finite supremum $\sup E_x$. Define $0 < \epsilon_x := \frac{1}{2} \sup E_x < \sup E_x$. Note the important thing that $\epsilon_x \in E_x$. Let us also observe that:

$$A \subseteq \bigcup_{x \in A} B_{\epsilon_x}(x) \subseteq \bigcup_{\alpha \in \mathcal{F}} \mathcal{O}_\alpha. \quad (2.2)$$

The first inclusion is obvious, while the second one follows from the above discussion.

We now need to prove a lemma:

Lemma 2.12. *If A is sequentially compact, then*

$$\inf_{x \in A} \epsilon_x =: 2\epsilon_0 > 0.$$

In other words, there exists $\epsilon_0 > 0$ such that $B_{\epsilon_0}(x) \subseteq B_{\epsilon_x}(x)$, for every $x \in A$.

Proof. Assume that $\inf_{x \in A} \epsilon_x = 0$. This implies that there exists a sequence $\{x_n\}_{n \geq 1} \subseteq A$ such that $\epsilon_{x_n} \leq 1/n$ for every $n \geq 1$. Since A is sequentially compact, there exists a convergent subsequence $\{x_{n_k}\}_{k \geq 1}$ which converges to a point $x_0 \in A$, i.e.

$$\lim_{k \rightarrow \infty} x_{n_k} = x_0. \quad (2.3)$$

Because x_0 belongs to A , we can find an open set $\mathcal{O}_{\alpha(x_0)}$ which contains x_0 , thus we can find $\epsilon_1 > 0$ such that

$$B_{\epsilon_1}(x_0) \subseteq \mathcal{O}_{\alpha(x_0)}. \quad (2.4)$$

Now (2.3) implies that there exists $K > 0$ large enough such that:

$$d(x_{n_k}, x_0) \leq \epsilon_1/4, \quad \text{whenever } k > K. \quad (2.5)$$

If y belongs to $B_{\epsilon_1/4}(x_{n_k})$ (i.e. $d(y, x_{n_k}) < \epsilon_1/4$), then the triangle inequality implies (use also (2.5)):

$$d(y, x_0) \leq d(y, x_{n_k}) + d(x_{n_k}, x_0) < \epsilon_1/2 < \epsilon_1, \quad k > K.$$

But this shows that we must have $y \in B_{\epsilon_1}(x_0)$, or:

$$B_{\epsilon_1/4}(x_{n_k}) \subseteq B_{\epsilon_1}(x_0) \subseteq \mathcal{O}_{\alpha(x_0)}, \quad \forall k > K. \quad (2.6)$$

Thus we got the inclusion

$$B_{\epsilon_1/4}(x_{n_k}) \subseteq \mathcal{O}_{\alpha(x_0)}, \quad \forall k > K,$$

which shows that $\epsilon_1/4$ must be less or equal than $2\epsilon_{x_{n_k}}$, or $\epsilon_1/8 \leq \epsilon_{x_{n_k}}$, for every $k > K$. But this is in contradiction with the fact that $\epsilon_{x_n} \leq 1/n$ for every $n \geq 1$. \square

Finishing the proof of Theorem 2.11. We now use Proposition 2.9, and find a finite ϵ_0 -net for A . Thus we can choose $\{y_1, \dots, y_N\} \subseteq A$ such that

$$A \subseteq \bigcup_{n=1}^N B_{\epsilon_0}(y_n) \subseteq \bigcup_{n=1}^N B_{\epsilon_{y_n}}(y_n) \subseteq \bigcup_{n=1}^N \mathcal{O}_n,$$

where \mathcal{O}_n is one of the possibly many open sets which contain $B_{\epsilon_{y_n}}(y_n)$. We have thus extracted our finite subcovering of A and the proof of the theorem is over. \square

2.3 The Heine-Borel Theorem

Theorem 2.13. *Consider \mathbb{R}^d with the Euclidian distance. In this metric space, a set A is (sequentially) compact if and only if A is both bounded and closed.*

Proof. We first show that a compact set is bounded and closed; this fact holds for all metric spaces, not just for the Euclidian ones. We already know that a compact set A is bounded (see Lemma 2.10). Let us prove that it is closed. Assume it is not. It means that the complementary A^c is not open, that is we can find a point $a \notin A$ which is not an interior point of A^c . Thus for every $n \geq 1$ we can find a point $x_n \in A$ such that $x_n \in B_{1/n}(a)$. The sequence $\{x_n\}_{n \geq 1} \subset A$ converges to a , thus any of its subsequences must converge to the same limit. But A is sequentially compact, thus there exists a subsequence $\{x_{n_k}\}_{k \geq 1}$ which converges to some point of A , which has to be a . This contradicts the fact that $a \notin A$.

Now let us show the other implication. Assume that A is bounded and consider an arbitrary sequence $\{x_n\}_{n \geq 1} \subset A$. The Bolzano-Weierstrass theorem implies the existence of a subsequence $\{x_{n_k}\}_{k \geq 1}$ which converges to some point $a \in \mathbb{R}^d$. Thus $a \in \overline{A}$, and due to Theorem 1.3 we know that $A = \overline{A}$, thus $a \in A$. This proves that A is sequentially compact, therefore compact. \square

3 Continuous functions on metric spaces

Let (X, d) and (Y, ρ) be two metric spaces. If $A \subset X$, the image of A through f is the set

$$f(A) := \{y \in Y : \text{there exists } x_y \in A \text{ such that } f(x_y) = y\} \subset Y.$$

If $B \subset Y$ the preimage of B through f is the set

$$f^{-1}(B) := \{x \in X : \text{such that } f(x) \in B\} \subset X.$$

Note that the notation $f^{-1}(B)$ does not imply that f is invertible.

Lemma 3.1. *If $A_1 \subset A_2 \subset X$ and $B_1 \subset B_2 \subset Y$ then $f(A_1) \subset f(A_2)$ and $f^{-1}(B_1) \subset f^{-1}(B_2)$.*

Proof. We only prove the first inclusion. Assume that $y \in f(A_1)$. Then there exists $x_y \in A_1$ such that $f(x_y) = y$. But at the same time $x_y \in A_2$, hence $y \in f(A_2)$. \square

A map $f : X \rightarrow Y$ is said to be continuous at a point $a \in X$ if for every $\epsilon > 0$ there exists $\delta > 0$ such that

$$B_\delta(a) \subset f^{-1}(B_\epsilon(f(a))), \quad (3.1)$$

which implies that $f(B_\delta(a)) \subset B_\epsilon(f(a))$. The function is continuous on X if it is continuous at all the points of X .

Theorem 3.2. *A function between two metric spaces $f : X \rightarrow Y$ is continuous on X if and only if for every nonempty open set $V \subset Y$ we have that $f^{-1}(V)$ is open in X .*

Proof. First we assume that f is continuous on X . Let V a nonempty open set in Y . If $f^{-1}(V)$ is empty then we know that it is open. Otherwise, let $a \in f^{-1}(V)$. Thus $f(a) \in V$. Since V is open, $f(a)$ is an interior point of V , thus there exists $\epsilon > 0$ such that $B_\epsilon(f(a)) \subset V$. Applying Lemma 3.1 we get that $f^{-1}(B_\epsilon(f(a))) \subset f^{-1}(V)$. But from (3.1) it follows that $B_\delta(a) \subset f^{-1}(V)$, thus a is an interior point.

We now assume that f returns any nonempty open set V of Y in an open set $f^{-1}(V)$ of X . Fix $a \in X$. Let $\epsilon > 0$ and consider the ball $B_\epsilon(f(a))$. Lemma 1.1 implies that $V = B_\epsilon(f(a))$ is open in Y . Thus $f^{-1}(B_\epsilon(f(a)))$ must be open in X . Since $a \in f^{-1}(B_\epsilon(f(a)))$, it must be an interior point. Thus there exists $\delta > 0$ such that $B_\delta(a) \subset f^{-1}(B_\epsilon(f(a)))$, which shows that f is continuous at a . \square

Let (X, d) and (Y, ρ) be two metric spaces and consider a subset $A \subset X$. We can organize A as a metric space with the natural distance d_A induced by d . We say that the map $f : A \mapsto Y$ is continuous on A if it is continuous between the metric spaces (A, d_A) and (Y, ρ) .

We say that $f : A \mapsto Y$ is sequentially continuous at a point $a \in A$ if for every sequence $\{x_n\}_{n \geq 1} \subset A$ which converges to a we have that $\{f(x_n)\}_{n \geq 1} \subset Y$ converges to $f(a)$. We say that $f : A \mapsto Y$ is sequentially continuous on A if it is sequentially continuous at all points of A .

Theorem 3.3. *With the above notation, consider a map $f : A \mapsto Y$. Then f is continuous on A if and only if it is sequentially continuous on A .*

Proof. First, assume that f is continuous at $a \in A$. Consider any sequence $\{x_n\}_{n \geq 1} \subset A$ which converges to a . From (3.1) we know that for every $\epsilon > 0$ we have that $\rho(f(x_n), f(a)) < \epsilon$ if $d(x_n, a) < \delta$. But the second inequality holds if n is larger than some $N_\delta \geq 1$. Thus $\{f(x_n)\}_{n \geq 1} \subset Y$ converges to $f(a)$.

Second, assume that f is sequentially continuous at $a \in A$. We will show that f must be continuous at a . Suppose this is not true: it means that there exists $\epsilon_0 > 0$ such that for all $\delta > 0$ we have that $B_\delta(a) \not\subset f^{-1}(B_{\epsilon_0}(f(a)))$. By letting $\delta = 1/n$ for all $n \geq 1$, we can find a point

$x_n \in B_{\frac{1}{n}}(a)$ such that $f(x_n) \notin B_{\epsilon_0}(f(a))$, or $\rho(f(x_n), f(a)) \geq \epsilon_0$. In this way we constructed a sequence $\{x_n\}_{n \geq 1} \subset A$ which converges to a while $\{f(x_n)\}_{n \geq 1}$ does not converge to $f(a)$, contradiction. \square

Theorem 3.4. *With the above notation, consider a continuous map $f : A \mapsto Y$ where $A \subset X$ is compact. Then $f(A)$ is compact.*

Proof. We show that $f(A)$ is sequentially compact. Let $\{y_n\}_{n \geq 1} \subset f(A)$ be an arbitrary sequence. There exists $\{x_n\}_{n \geq 1} \subset A$ such that $f(x_n) = y_n$. Since A is sequentially compact, there exists a subsequence $\{x_{n_k}\}_{k \geq 1} \subset \{x_n\}_{n \geq 1}$ which converges to some point $a \in A$. But f is sequentially continuous at a , hence $y_{n_k} = f(x_{n_k})$ converges to $f(a) \in f(A)$. Hence $f(A)$ is sequentially compact. \square

The next lemma recalls a general result which says that real continuous functions defined on compact sets attain their extremal values. See also Theorem 10.61 in Wade.

Lemma 3.5. *Let (X, d) be a metric space and let $H \subset X$ be a compact set. Let $f : H \mapsto \mathbb{R}$ be continuous on H . Then there exist x_m and x_M in H such that $f(x_M) = \sup_{x \in H} f(x)$ and $f(x_m) = \inf_{x \in H} f(x)$.*

Proof. We only prove this for $\sup_{x \in H} f(x)$. Let $B := f(H) \subset \mathbb{R}$. Let us show that there exists a sequence $\{x_n\}_{n \geq 1} \subset H$ such that $\lim_{n \rightarrow \infty} f(x_n) = \sup_{x \in H} f(x) = \sup(B)$.

Since B is compact, it is bounded. Thus $\sup(B) = \sup_{x \in H} f(x) < \infty$. For every $n \geq 1$ we know that $\sup(B) - 1/n$ is not an upper bound for B , thus there must exist $x_n \in H$ such that $\sup(B) - 1/n < f(x_n) \leq \sup(B)$. Thus $\lim_{n \rightarrow \infty} f(x_n) = \sup(B)$.

Because H is compact, we can find a subsequence $\{x_{n_k}\}_{k \geq 1}$ which converges towards some point $a \in H$. Since f is continuous, we have that $\lim_{k \rightarrow \infty} f(x_{n_k}) = f(a)$. Since $\{f(x_{n_k})\}_{k \geq 1}$ is a subsequence of the convergent sequence $\{f(x_n)\}_{n \geq 1}$, we must have $f(a) = \sup(B)$. Thus we can choose x_M to be a . \square

We say that $f : A \mapsto Y$ is uniformly continuous on A if for every $\epsilon > 0$ there exists $\delta > 0$ such that $\rho(f(x), f(y)) < \epsilon$ as soon as $x, y \in A$ and $d(x, y) < \delta$. Clearly, if f is uniformly continuous on A then it is also continuous. The next result gives sufficient conditions for the reciprocal statement:

Lemma 3.6. *Let (X, d) and (Y, ρ) be two metric spaces and let $H \subset X$ be a compact set. Let $f : H \mapsto Y$ be continuous on H . Then f is uniformly continuous on H .*

Proof. Assume that the conclusion is false. Then there exists $\epsilon_0 > 0$ such that regardless how large $n \geq 1$ is, we may find two points x_n and y_n in H which obey $d(x_n, y_n) < \frac{1}{n}$ and $\rho(f(x_n), f(y_n)) \geq \epsilon_0$. Since H is sequentially compact, there exists a subsequence $\{x_{n_k}\}_{k \geq 1}$ which converges to some point $a \in H$. Because $d(y_{n_k}, a) \leq \frac{1}{k} + d(x_{n_k}, a)$ for all $k \geq 1$, it follows that y_{n_k} also converges to a . The function f is sequentially compact at a , thus both $f(x_{n_k})$ and $f(y_{n_k})$ converge to $f(a)$. In particular, this contradicts our assumption that $\rho(f(x_{n_k}), f(y_{n_k})) \geq \epsilon_0$ for all k . \square

4 Banach's fixed point theorem

Definition 4.1. Let (X, d) be a metric space. A map $F : X \rightarrow X$ is called a contraction if there exists $\alpha \in [0, 1)$ such that:

$$d(F(x), F(y)) \leq \alpha d(x, y), \quad \forall x, y \in X. \quad (4.2)$$

A point $x \in X$ is a fixed point for F if $F(x) = x$.

Theorem 4.2. Let (X, d) be a complete metric space and $F : X \rightarrow X$ a contraction. Then F has a unique fixed point.

Proof. We start by showing uniqueness. Assume that there exist $a, b \in X$ such that $F(a) = a$ and $F(b) = b$. Then (4.2) implies that

$$0 \leq d(a, b) = d(F(a), F(b)) \leq \alpha d(a, b), \quad (1 - \alpha)d(a, b) \leq 0,$$

i.e. $d(a, b) = 0$ and $a = b$.

Now let us construct such a fixed point. Consider the sequence $\{y_n\}_{n \geq 1} \subset X$, where y_1 is arbitrary and $y_n := F(y_{n-1})$ for every $n \geq 2$. We will show two things:

(i). The sequence is Cauchy in X , thus convergent to a limit y_∞ because we assumed X to be complete;

(ii). y_∞ is a fixed point for F .

Let us start with (i). For every $\epsilon > 0$ we will construct $N(\epsilon) > 0$ such that for all $p \geq q \geq N(\epsilon)$ we have $d(y_q, y_p) < \epsilon$. In other words:

$$d(y_q, y_{q+k}) < \epsilon, \quad \forall k \geq 0, \quad \forall q \geq N(\epsilon). \quad (4.3)$$

If $k \geq 1$, the triangle inequality implies:

$$\begin{aligned} d(y_q, y_{q+k}) &\leq d(y_q, y_{q+1}) + d(y_{q+1}, y_{q+k}) \\ &\leq d(y_q, y_{q+1}) + d(y_{q+1}, y_{q+2}) + d(y_{q+2}, y_{q+k}) \\ &\leq \sum_{i=0}^{k-1} d(y_{q+i}, y_{q+i+1}). \end{aligned} \quad (4.4)$$

For every $n \geq 1$ we have:

$$d(y_n, y_{n+1}) = d(F(y_{n-1}), F(y_n)) \leq \alpha d(y_{n-1}, y_n) \leq \dots \leq \alpha^{n-1} d(y_1, y_2), \quad \forall n \geq 1.$$

Thus $d(y_{q+i}, y_{q+i+1}) \leq \alpha^{q+i-1} d(y_1, y_2)$ for all $q \geq 1$ and $i \geq 0$. Together with (4.4), this implies:

$$d(y_q, y_{q+k}) \leq \alpha^{q-1} d(y_1, y_2) (1 + \dots + \alpha^{k-1}) \leq \frac{\alpha^{q-1}}{1 - \alpha} d(y_1, y_2), \quad \forall k \geq 1.$$

Because $\alpha < 1$, then $\lim_{q \rightarrow \infty} \alpha^q = 0$ and (4.3) follows. We conclude that there exists $y_\infty \in X$ such that

$$\lim_{n \rightarrow \infty} d(y_n, y_\infty) = 0. \quad (4.5)$$

Now we prove (ii). For every $n \geq 1$ we have:

$$d(F(y_\infty), y_\infty) \leq d(F(y_\infty), F(y_n)) + d(F(y_n), y_\infty).$$

But $d(F(y_\infty), F(y_n)) \leq \alpha d(y_\infty, y_n) \rightarrow 0$ and $d(F(y_n), y_\infty) = d(y_{n+1}, y_\infty) \rightarrow 0$ when $n \rightarrow \infty$, thus $d(F(y_\infty), y_\infty) = 0$ and $F(y_\infty) = y_\infty$. \square

5 Local existence and uniqueness for first order ODE's

We start with some general facts about functional spaces.

5.1 Spaces of bounded/continuous functions

Proposition 5.1. *Let (A, d) be a metric space, $(Y, \|\cdot\|)$ a normed space, and H an arbitrary non-empty subset of A . We define*

$$B(H; Y) := \{f : H \rightarrow Y : \sup_{x \in H} \|f(x)\| < \infty\}.$$

Define the map $\|\cdot\|_\infty : B(H; Y) \rightarrow \mathbb{R}_+$, $\|f\|_\infty := \sup_{x \in H} \|f(x)\|$. Then the space $(B(H; Y), \|\cdot\|_\infty)$ is a normed space, and the map $d_\infty(f, g) := \|f - g\|_\infty$ defines a metric.

Proof. Clearly, $\|f\|_\infty = 0$ if and only if $f(x) = 0$ for all $x \in H$. Moreover,

$$\|\lambda f\|_\infty = \sup_{x \in H} \|\lambda f(x)\| = |\lambda| \sup_{x \in H} \|f(x)\| = |\lambda| \|f\|_\infty.$$

Finally, let us prove the triangle inequality. Take $f, g \in B(H; Y)$; then for every $x \in H$ we apply the triangle inequality in $(Y, \|\cdot\|)$:

$$\|f(x) + g(x)\| \leq \|f(x)\| + \|g(x)\| \leq \|f\|_\infty + \|g\|_\infty.$$

Thus $\|f\|_\infty + \|g\|_\infty$ is an upper bound for the set $\{\|f(x) + g(x)\| : x \in H\}$, hence

$$\sup_{x \in H} \|f(x) + g(x)\| = \|f + g\|_\infty \leq \|f\|_\infty + \|g\|_\infty.$$

Note that $d_\infty(f, g) := \|f - g\|_\infty$ is the metric induced by the norm. □

Proposition 5.2. *Denote by $C(H; Y)$ the subset of $B(H; Y)$ where the functions are also continuous. Assume that $(Y, \|\cdot\|)$ is a Banach space (a complete normed space). Then $(C(H; Y), \|\cdot\|_\infty)$ is a Banach space, too.*

Proof. We need to prove that every Cauchy sequence is convergent. Assume that $\{f_n\}_{n \geq 1} \subset C(H; Y)$ is Cauchy, i.e. for every $\epsilon > 0$ one can find $N_C(\epsilon) > 0$ such that $\|f_p - f_q\|_\infty < \epsilon$ if $p, q > N_C(\epsilon)$. We have to show that the sequence has a limit f which belongs to $C(H; Y)$.

We first construct f . For every $x_0 \in H$ we consider the sequence $\{f_n(x_0)\}_{n \geq 1} \subset Y$. Note the difference between $\{f_n(x_0)\}_{n \geq 1}$ (a sequence of vectors from Y) and $\{f_n\}_{n \geq 1}$ (a sequence of functions from $C(H; Y)$). It is easy to see that $\{f_n(x_0)\}_{n \geq 1}$ is Cauchy in Y (exercise), and because Y is complete, then $\{f_n(x_0)\}_{n \geq 1}$ has a limit in Y . We denote it with $f(x_0)$. Moreover, since $\{f_n\}_{n \geq 1}$ is Cauchy it must be bounded, i.e. $\|f_n\|_\infty \leq M < \infty$ for all $n \geq 1$. Thus we have:

$$\|f(x)\| = \lim_{n \rightarrow \infty} \|f_n(x)\| \leq M, \quad \forall x \in H,$$

therefore $\|f\|_\infty < \infty$.

Second, we prove the "uniform convergence" part, or the convergence in the norm $\|\cdot\|_\infty$. More precisely, it means that for every $\epsilon > 0$ we must construct $N_1(\epsilon) > 0$ so that:

$$\sup_{x \in H} \|f(x) - f_n(x)\| < \epsilon \quad \text{whenever } n > N_1(\epsilon). \quad (5.1)$$

In order to do that, take an arbitrary point $x \in H$. For every $p, n \geq 1$ we have

$$\begin{aligned} \|f(x) - f_n(x)\| &\leq \|f(x) - f_p(x)\| + \|f_p(x) - f_n(x)\| \\ &\leq \|f(x) - f_p(x)\| + \|f_p - f_n\|_\infty. \end{aligned} \quad (5.2)$$

If we choose $n, p > N_C(\epsilon/2)$, then we have $\|f_p - f_n\|_\infty < \epsilon/2$ and

$$\|f(x) - f_n(x)\| \leq \|f(x) - f_p(x)\| + \epsilon/2, \quad n, p > N_C(\epsilon/2).$$

But the above left hand side does not depend on p , thus if we take $p \rightarrow \infty$ on the right hand side, we get:

$$\|f(x) - f_n(x)\| \leq \epsilon/2 < \epsilon, \quad n > N_C(\epsilon/2). \quad (5.3)$$

Note that this inequality holds true *for every* x . This means that $\epsilon/2$ is an upper bound for the set $\{\|f(x) - f_n(x)\| : x \in H\}$, hence (5.1) holds true with $N_1(\epsilon) = N_C(\epsilon/2)$.

Third, we must prove that f is a continuous function on H . Fix some point $a \in H$. Choose $\epsilon > 0$. Since $\lim_{n \rightarrow \infty} f_n(a) = f(a)$, we can find $N_2(\epsilon, a) > 0$ such that $\|f_n(a) - f(a)\| < \epsilon$ whenever $n > N_2$. We define $n_1 := \max\{N_1(\epsilon/3), N_2(\epsilon/3, a)\}$. Because f_{n_1} is continuous at a , we can find $\delta(\epsilon, a) > 0$ so that for every $x \in H$ with $d(x, a) < \delta$ we have $\|f_{n_1}(x) - f_{n_1}(a)\| < \epsilon/3$. Thus if $x \in H$ with $d(x, a) < \delta$ we have:

$$\begin{aligned} \|f(x) - f(a)\| &\leq \|f(x) - f_{n_1}(x)\| + \|f_{n_1}(x) - f_{n_1}(a)\| + \|f_{n_1}(a) - f(a)\| \\ &< 2\|f - f_{n_1}\|_\infty + \|f_{n_1}(x) - f_{n_1}(a)\| < \epsilon. \end{aligned} \quad (5.4)$$

Since a is arbitrary, we can conclude that f is continuous on H , thus belongs to $C(H; Y)$. Therefore we can rewrite (5.1) as:

$$\|f - f_n\|_\infty < \epsilon \quad \text{whenever} \quad n > N_1(\epsilon), \quad (5.5)$$

and the proof is over. \square

Remark 5.3. The "ordinary" convergence in the functional space $(C(H; Y), \|\cdot\|_\infty)$ (given in (5.5)) is nothing but the uniform convergence of a sequence of functions defined on the set H (see (5.1)). One can find more details in Wade, exercise 10.6.6 in Chapter 10.6 (page 376).

5.2 The main theorem

Let U be an open set in \mathbb{R}^d , $d \geq 1$, and $I \subset \mathbb{R}$ an open interval. Assume that there exist $\mathbf{y}_0 \in U$ and $r_0, \delta_0 > 0$ such that $\overline{B_{r_0}(\mathbf{y}_0)} \subset U$ and $[t_0 - \delta_0, t_0 + \delta_0] \subset I$.

We consider a continuous function $\mathbf{f} : I \times U \rightarrow \mathbb{R}^d$ for which there exists $L > 0$ such that

$$\|\mathbf{f}(t, \mathbf{x}) - \mathbf{f}(t, \mathbf{y})\| \leq L\|\mathbf{x} - \mathbf{y}\|, \quad \forall t \in [t_0 - \delta_0, t_0 + \delta_0], \quad \forall \mathbf{x}, \mathbf{y} \in \overline{B_{r_0}(\mathbf{y}_0)}. \quad (5.6)$$

We define the compact set $H_0 := [t_0 - \delta_0, t_0 + \delta_0] \times \overline{B_{r_0}(\mathbf{y}_0)} \subset \mathbb{R}^{d+1}$. Because \mathbf{f} is continuous, the same is true for $\|\mathbf{f}\|$, thus according to Lemma 3.5 we can find $M < \infty$ such that

$$\sup_{[t, \mathbf{x}] \in H_0} \|\mathbf{f}(t, \mathbf{x})\| =: M < \infty. \quad (5.7)$$

Consider the initial value problem:

$$\mathbf{y}'(t) = \mathbf{f}(t, \mathbf{y}(t)), \quad \mathbf{y}(t_0) = \mathbf{y}_0. \quad (5.8)$$

Theorem 5.4. Define

$$\delta_1 := \min\{\delta_0, r_0/M, 1/L\}.$$

Then the equation (5.8) has a unique solution for $t \in]t_0 - \delta_1, t_0 + \delta_1[$.

Proof. Take some $0 < \delta < \delta_1$ and define the compact interval $K := [t_0 - \delta, t_0 + \delta] \subset \mathbb{R}$. Then any continuous function $\phi : K \rightarrow \mathbb{R}^d$ is automatically bounded, and since the Euclidian space $Y = \mathbb{R}^d$ is a Banach space, we can conclude from Proposition 5.2 that the space $(C(K; \mathbb{R}^d), d_\infty)$ of continuous functions defined on the compact K with values in \mathbb{R}^d is a complete metric space.

Define

$$X := \{g \in C(K; \mathbb{R}^d) : g(t) \in \overline{B_{r_0}(\mathbf{y}_0)}, \forall t \in K\}. \quad (5.9)$$

Lemma 5.5. The metric space (X, d_∞) is complete.

Proof. Consider a Cauchy sequence $\{f_n\}_{n \geq 1} \subset X$. Because $(C(K; \mathbb{R}^d), d_\infty)$ is complete, we can find $f_\infty \in C(K; \mathbb{R}^d)$ such that $\lim_{n \rightarrow \infty} d_\infty(f_n, f_\infty) = 0$. Thus for every $t \in K$ we have

$$f_\infty(t) = \lim_{n \rightarrow \infty} f_n(t), \quad \lim_{n \rightarrow \infty} \|f_n(t) - f_\infty(t)\| = 0.$$

Since by assumption $\|f_n(t) - \mathbf{y}_0\| \leq r_0$ for all t and n , we have

$$\|f_\infty(t) - \mathbf{y}_0\| = \lim_{n \rightarrow \infty} \|f_n(t) - \mathbf{y}_0\| \leq r_0, \quad \forall t \in K,$$

which implies that $f_\infty \in X$. □

Lemma 5.6. Define the map $F : X \rightarrow C(K; \mathbb{R}^d)$

$$[F(g)](t) := \mathbf{y}_0 + \int_{t_0}^t \mathbf{f}(s, g(s)) ds, \quad \forall t \in K,$$

where \mathbf{f} is given in (5.6). Then:

- (i). The range of F belongs to X ;
- (ii). $F : X \rightarrow X$ is a contraction.

Proof. (i). Because $g(s) \in \overline{B_{r_0}(\mathbf{y}_0)}$ for all $s \in K$, we have that $(s, g(s)) \in H_0$ for all $s \in K$. Thus (see (5.7)) $\sup_{s \in K} \|\mathbf{f}(s, g(s))\| \leq M$ and

$$\|[F(g)](t) - \mathbf{y}_0\| \leq \left\| \int_{t_0}^t \mathbf{f}(s, g(s)) ds \right\| \leq M\delta < r_0, \quad \forall t \in K,$$

which means that $[F(g)](t) \in \overline{B_{r_0}(\mathbf{y}_0)}$ for all $t \in K$.

- (ii). Consider two functions $g, h \in X$. We have

$$d_\infty(F(g), F(h)) = \sup_{t \in K} \|[F(g)](t) - [F(h)](t)\|.$$

But the Lipschitz condition from (5.6) implies:

$$\begin{aligned} |[F(g)](t) - [F(h)](t)| &= \left| \int_{t_0}^t [\mathbf{f}(s, g(s)) - \mathbf{f}(s, h(s))] ds \right| \leq (\delta L) \sup_{s \in K} \|g(s) - h(s)\| \\ &\leq (\delta L) d_\infty(g, h), \quad \forall t \in K. \end{aligned} \tag{5.10}$$

It means that $d_\infty(F(g), F(h)) \leq (\delta L) d_\infty(g, h)$ for all $g, h \in X$, and remember that $\delta L < 1$. Thus F is a contraction. □

Finishing the proof of Theorem 5.4. We have seen that F is a contraction on X . Then Theorem 4.2 implies that there exists a continuous function $\mathbf{y} : K \rightarrow \overline{B_{r_0}(\mathbf{y}_0)}$ such that

$$\mathbf{y}(t) = [F(\mathbf{y})](t) = \mathbf{y}_0 + \int_{t_0}^t \mathbf{f}(s, \mathbf{y}(s)) ds, \quad t \in [t_0 - \delta, t_0 + \delta].$$

It means that \mathbf{y} is differentiable for $t \in]t_0 - \delta, t_0 + \delta[$ and (5.8) is satisfied. □

Remark 5.7. Choose $0 < \delta < \delta_1$. Define the sequence of functions $\mathbf{y}_k : [t_0 - \delta, t_0 + \delta] \rightarrow \mathbb{R}^d$, $k \geq 1$, where $\mathbf{y}_1(t) = \mathbf{y}_0$ and

$$\mathbf{y}_{k+1}(t) = \mathbf{y}_0 + \int_{t_0}^t \mathbf{f}(s, \mathbf{y}_k(s)) ds, \quad k \geq 1.$$

We see that $\mathbf{y}_{k+1} = F(\mathbf{y}_k)$, where F is given by Lemma 5.6. A direct use of Lemma 5.6 (ii) implies that $\{\mathbf{y}_k\}_{k \geq 1}$ converges uniformly on the interval $[t_0 - \delta, t_0 + \delta]$ towards a continuous function \mathbf{y}_∞ which obeys the fixed point equation

$$\mathbf{y}_\infty(t) = \mathbf{y}_0 + \int_{t_0}^t \mathbf{f}(s, \mathbf{y}_\infty(s)) ds,$$

thus solving (5.8). This is Picard's iteration method.

6 The inverse function theorem

In this section we only work with the Euclidian space \mathbb{R}^d , whose norm is defined by $\|\mathbf{x}\| = \sqrt{\sum_{j=1}^d |x_j|^2}$.

Lemma 6.1. *Let A be a $d \times d$ matrix with real components $\{a_{jk}\}$. Define the quantity $\|A\|_{\text{HS}} := \sqrt{\sum_{j=1}^d \sum_{k=1}^d |a_{jk}|^2}$. Then*

$$\|A\mathbf{x}\| \leq \|A\|_{\text{HS}} \|\mathbf{x}\|, \quad \forall \mathbf{x} \in \mathbb{R}^d. \quad (6.1)$$

Proof. From the Cauchy-Schwarz inequality we have:

$$|(A\mathbf{x})_j|^2 = \left(\sum_{k=1}^d a_{jk} x_k \right)^2 \leq \sum_{m=1}^d |a_{jm}|^2 \sum_{n=1}^d |x_n|^2 = \sum_{m=1}^d |a_{jm}|^2 \|\mathbf{x}\|^2,$$

and the lemma follows after summation with respect to j . \square

Lemma 6.2. *Let $K := \overline{B_\delta(\mathbf{x}_0)} = \{\mathbf{y} \in \mathbb{R}^d : \|\mathbf{y} - \mathbf{x}_0\| \leq \delta\}$ be a closed ball in \mathbb{R}^d . Let $\phi : \mathbb{R}^d \mapsto \mathbb{R}$ be a $C^1(K)$ map (which means that $\partial_j \phi$ exist for all j and are continuous functions on K). Denote by $\|\partial_j \phi\|_\infty = \sup_{\mathbf{x} \in K} |\partial_j \phi(\mathbf{x})| < \infty$. Then for every $\mathbf{u}, \mathbf{w} \in K$ we have:*

$$|\phi(\mathbf{u}) - \phi(\mathbf{w})| \leq \sqrt{\sum_{j=1}^d \|\partial_j \phi\|_\infty^2} \|\mathbf{u} - \mathbf{w}\|. \quad (6.2)$$

Proof. Define the real valued map $f(t) = \phi((1-t)\mathbf{w} + t\mathbf{u})$, $0 \leq t \leq 1$. Applying the chain rule we obtain:

$$f'(t) = \sum_{j=1}^d (u_j - w_j) (\partial_j \phi)((1-t)\mathbf{w} + t\mathbf{u}),$$

thus the Cauchy-Schwarz inequality implies:

$$|f'(t)| \leq \sqrt{\sum_{j=1}^d |\partial_j \phi((1-t)\mathbf{w} + t\mathbf{u})|^2} \|\mathbf{u} - \mathbf{w}\| \leq \sqrt{\sum_{j=1}^d \|\partial_j \phi\|_\infty^2} \|\mathbf{u} - \mathbf{w}\|, \quad \forall 0 < t < 1.$$

Since $\phi(\mathbf{u}) - \phi(\mathbf{w}) = f(1) - f(0) = \int_0^1 f'(t) dt$, we obtain:

$$|\phi(\mathbf{u}) - \phi(\mathbf{w})| \leq \int_0^1 |f'(t)| dt \leq \sqrt{\sum_{j=1}^d \|\partial_j \phi\|_\infty^2} \|\mathbf{u} - \mathbf{w}\|$$

which proves (6.2). \square

Lemma 6.3. *Let K be as above. Let $\mathbf{f} : \mathbb{R}^d \mapsto \mathbb{R}^d$ a vector valued map which is $C^1(K)$ (which means that $\partial_j f_k$ exist and are continuous functions on K). Define*

$$\|\Delta \mathbf{f}\|_{\infty, K} := \sqrt{\sum_{k=1}^d \sum_{j=1}^d \|\partial_j f_k\|_\infty^2}.$$

Then we have:

$$\|\mathbf{f}(\mathbf{u}) - \mathbf{f}(\mathbf{w})\| \leq \|\Delta \mathbf{f}\|_{\infty, K} \|\mathbf{u} - \mathbf{w}\|, \quad \forall \mathbf{u}, \mathbf{w} \in K. \quad (6.3)$$

Proof. Use (6.2) with ϕ replaced by f_k . We have:

$$|f_k(\mathbf{u}) - f_k(\mathbf{w})|^2 \leq \sum_{j=1}^d \|\partial_j f_k\|_\infty^2 \|\mathbf{u} - \mathbf{w}\|^2$$

and then sum over k . □

Lemma 6.4. *Using the above notation, define $\mathbf{g}(\mathbf{x}) = \mathbf{f}(\mathbf{x}) - [D\mathbf{f}(\mathbf{x}_0)]\mathbf{x}$, where $[D\mathbf{f}(\mathbf{x}_0)]$ is the Jacobi matrix with elements $[D\mathbf{f}(\mathbf{x}_0)]_{kj} = (\partial_j f_k)(\mathbf{x}_0)$. Then for every $\beta > 0$ there exists a $\delta_\beta > 0$ such that for every $0 < \delta < \delta_\beta$ we have:*

$$\|\mathbf{g}(\mathbf{u}) - \mathbf{g}(\mathbf{w})\| \leq \beta \|\mathbf{u} - \mathbf{w}\|, \quad \forall \mathbf{u}, \mathbf{w} \in K. \quad (6.4)$$

Proof. A straightforward computation gives $\partial_j g_k(\mathbf{x}) = \partial_j f_k(\mathbf{x}) - \partial_j f_k(\mathbf{x}_0)$. Thus $\|\partial_j g_k\|_\infty$ can be made arbitrarily small when δ gets smaller, because \mathbf{f} has continuous partial derivatives. It follows that $\|\Delta \mathbf{g}\|_{\infty, K} \leq \beta$ whenever δ gets smaller than some small enough δ_β , and then we can use (6.3) with \mathbf{g} instead of \mathbf{f} . □

Lemma 6.5. *Let $\mathbf{a} \in \mathbb{R}^d$ and let $U \subset \mathbb{R}^d$ be an open set with $\mathbf{a} \in U$. Let \mathbf{f} be a $C^1(U)$ vector valued function, such that $[D\mathbf{f}(\mathbf{a})]$ is an invertible matrix. Then there exists $r > 0$ small enough such that the restriction of \mathbf{f} to $B_r(\mathbf{a})$ is injective.*

Proof. Assume the contrary: there exists a sequence $\{r_n\}_{n \geq 1}$ converging to zero such that for every $n \geq 1$ we can find two different points $\mathbf{x}_n \neq \mathbf{y}_n$ in $B_{r_n}(\mathbf{a})$ such that $\mathbf{f}(\mathbf{x}_n) = \mathbf{f}(\mathbf{y}_n)$. Define $\mathbf{g}(\mathbf{x}) = \mathbf{f}(\mathbf{x}) - [D\mathbf{f}(\mathbf{a})]\mathbf{x}$ on $B_{r_n}(\mathbf{a})$. Then we have $\mathbf{g}(\mathbf{x}_n) - \mathbf{g}(\mathbf{y}_n) = [D\mathbf{f}(\mathbf{a})](\mathbf{y}_n - \mathbf{x}_n)$ or:

$$\mathbf{y}_n - \mathbf{x}_n = [D\mathbf{f}(\mathbf{a})]^{-1}(\mathbf{g}(\mathbf{x}_n) - \mathbf{g}(\mathbf{y}_n)), \quad \forall n \geq 1.$$

Now using (6.1) we have:

$$\|\mathbf{y}_n - \mathbf{x}_n\| = \|[D\mathbf{f}(\mathbf{a})]^{-1}\|_{\text{HS}} \|\mathbf{g}(\mathbf{x}_n) - \mathbf{g}(\mathbf{y}_n)\|, \quad \forall n \geq 1.$$

Choosing $\beta = \frac{1}{1 + \|[D\mathbf{f}(\mathbf{a})]^{-1}\|_{\text{HS}}}$, then (6.4) claims that there exists some $r_\beta > 0$ sufficiently small such that for every $r_n \leq r_\beta$ we have $\|\mathbf{g}(\mathbf{x}_n) - \mathbf{g}(\mathbf{y}_n)\| \leq \beta \|\mathbf{y}_n - \mathbf{x}_n\|$. It follows that:

$$\|\mathbf{y}_n - \mathbf{x}_n\| \leq \frac{\|[D\mathbf{f}(\mathbf{a})]^{-1}\|_{\text{HS}}}{1 + \|[D\mathbf{f}(\mathbf{a})]^{-1}\|_{\text{HS}}} \|\mathbf{y}_n - \mathbf{x}_n\| < \|\mathbf{y}_n - \mathbf{x}_n\|, \quad \forall 0 < r_n < r_\beta,$$

which contradicts the assumption $\|\mathbf{y}_n - \mathbf{x}_n\| \neq 0$. □

Lemma 6.6. *Let \mathbf{f} be as in Lemma 6.5, and consider the injective restriction of f to $B_r(\mathbf{a})$. Then by eventually making r even smaller we have that the Jacobi matrix $[D\mathbf{f}(\mathbf{x})]$ is invertible for every $\mathbf{x} \in B_r(\mathbf{a})$.*

Proof. The matrix $[D\mathbf{f}(\mathbf{x})]$ is invertible if and only if its determinant $\det[D\mathbf{f}(\mathbf{x})] \neq 0$. But the determinant is a continuous function of \mathbf{x} since \mathbf{f} is C^1 . Because $[D\mathbf{f}(\mathbf{a})]$ is invertible, it follows that $|\det[D\mathbf{f}(\mathbf{a})]| > 0$. Being continuous at \mathbf{a} , the determinant has the property that $|\det[D\mathbf{f}(\mathbf{x})]| \geq |\det[D\mathbf{f}(\mathbf{a})]|/2 > 0$ on a small ball around \mathbf{a} . Thus $[D\mathbf{f}(\mathbf{x})]$ is invertible there. □

Theorem 6.7. *Let \mathbf{f} be C^1 on an open set containing $\mathbf{a} \in \mathbb{R}^d$, such that $[D\mathbf{f}(\mathbf{a})]$ is invertible. Then there exists $r > 0$ small enough such that the restriction of f to $B_r(\mathbf{a})$ is injective, and $[D\mathbf{f}(\mathbf{x})]$ is invertible for every $\mathbf{x} \in B_r(\mathbf{a})$. Moreover, if $V := f(B_r(\mathbf{a}))$, the following facts hold true:*

- (i). *The set V is open in \mathbb{R}^d ;*
- (ii). *The inverse $\mathbf{f}^{-1} : V \mapsto B_r(\mathbf{a})$ is a $C^1(V)$ function, and we have:*

$$[D\mathbf{f}^{-1}(\mathbf{y})] = [D\mathbf{f}(\mathbf{f}^{-1}(\mathbf{y}))]^{-1}.$$

Proof. The ball $B_r(\mathbf{a})$ has already been constructed in Lemma 6.6, hence we only need to prove (i) and (ii).

We start with (i). Assume that $\mathbf{y}_0 \in V$, thus equal to $\mathbf{f}(\mathbf{x}_0)$ for some $\mathbf{x}_0 \in B_r(\mathbf{a})$. We will show that \mathbf{y}_0 is an interior point of V . This means that we must show the existence of a small ball $B_\epsilon(\mathbf{y}_0)$ which is completely contained in V . In other words, we have to prove that there exists a sufficiently small $\epsilon > 0$ such that for every $\mathbf{y} \in \mathbb{R}^d$ with $\|\mathbf{y} - \mathbf{y}_0\| < \epsilon$ we can find a point $\mathbf{x}_y \in B_r(\mathbf{a})$ such that $\mathbf{f}(\mathbf{x}_y) = \mathbf{y} \in V$, hence $B_\epsilon(\mathbf{y}_0) \subset V$.

So the main question we need to answer is the solvability of the equation $\mathbf{f}(\mathbf{x}) = \mathbf{y}$. This equation is equivalent with:

$$0 = \mathbf{f}(\mathbf{x}) - \mathbf{y} = \mathbf{f}(\mathbf{x}) - \mathbf{f}(\mathbf{x}_0) - [D\mathbf{f}(\mathbf{x}_0)](\mathbf{x} - \mathbf{x}_0) + [D\mathbf{f}(\mathbf{x}_0)](\mathbf{x} - \mathbf{x}_0) + \mathbf{y}_0 - \mathbf{y}.$$

Denote by $\mathbf{g}(\mathbf{x}) = \mathbf{f}(\mathbf{x}) - [D\mathbf{f}(\mathbf{x}_0)]\mathbf{x}$. Then we have the equivalence between $0 = \mathbf{f}(\mathbf{x}) - \mathbf{y}$ and the equation

$$\mathbf{g}(\mathbf{x}) - \mathbf{g}(\mathbf{x}_0) + [D\mathbf{f}(\mathbf{x}_0)](\mathbf{x} - \mathbf{x}_0) + \mathbf{y}_0 - \mathbf{y} = 0.$$

Since $[D\mathbf{f}(\mathbf{x}_0)]$ is invertible, we can isolate \mathbf{x} and write another equivalent equation:

$$\mathbf{x} = \mathbf{x}_0 + [D\mathbf{f}(\mathbf{x}_0)]^{-1}(\mathbf{y} - \mathbf{y}_0) - [D\mathbf{f}(\mathbf{x}_0)]^{-1}(\mathbf{g}(\mathbf{x}) - \mathbf{g}(\mathbf{x}_0)).$$

This looks like a fixed point equation. Indeed, let us denote by

$$F_{\mathbf{y}}(\mathbf{x}) := \mathbf{x}_0 + [D\mathbf{f}(\mathbf{x}_0)]^{-1}(\mathbf{y} - \mathbf{y}_0) - [D\mathbf{f}(\mathbf{x}_0)]^{-1}(\mathbf{g}(\mathbf{x}) - \mathbf{g}(\mathbf{x}_0)). \quad (6.5)$$

It follows that if we can find a fixed point for $F_{\mathbf{y}}$, it will also solve the equation $\mathbf{f}(\mathbf{x}) = \mathbf{y}$.

We note first that using (6.1) we have:

$$\|F_{\mathbf{y}}(\mathbf{x}) - \mathbf{x}_0\| \leq \| [D\mathbf{f}(\mathbf{x}_0)]^{-1} \|_{\text{HS}} (\|\mathbf{y} - \mathbf{y}_0\| + \|\mathbf{g}(\mathbf{x}) - \mathbf{g}(\mathbf{x}_0)\|). \quad (6.6)$$

Choosing $\beta = \beta_1 = \frac{1}{3(1 + \| [D\mathbf{f}(\mathbf{x}_0)]^{-1} \|_{\text{HS}})}$ in (6.4), it follows that there exists a $\delta_1 > 0$ small enough, in any case smaller than $r - \|\mathbf{x}_0 - \mathbf{a}\|$, such that for every $\delta < \delta_1$ and $\|\mathbf{x} - \mathbf{x}_0\| < \delta$, we have $\|\mathbf{g}(\mathbf{x}) - \mathbf{g}(\mathbf{x}_0)\| \leq \beta_1 \|\mathbf{x} - \mathbf{x}_0\|$, thus:

$$\|F_{\mathbf{y}}(\mathbf{x}) - \mathbf{x}_0\| \leq \| [D\mathbf{f}(\mathbf{x}_0)]^{-1} \|_{\text{HS}} \|\mathbf{y} - \mathbf{y}_0\| + \frac{\delta}{3}, \quad \forall \mathbf{x} \in \overline{B_\delta(\mathbf{x}_0)} \subset B_r(\mathbf{a}). \quad (6.7)$$

In particular, if

$$\|\mathbf{y} - \mathbf{y}_0\| < \frac{\delta}{3(1 + \| [D\mathbf{f}(\mathbf{x}_0)]^{-1} \|_{\text{HS}})} =: \epsilon_\delta \quad (6.8)$$

then (6.7) states that

$$\|F_{\mathbf{y}}(\mathbf{x}) - \mathbf{x}_0\| \leq \frac{2\delta}{3} \leq \delta, \quad \forall \mathbf{x} \in \overline{B_\delta(\mathbf{x}_0)} \subset B_r(\mathbf{a}). \quad (6.9)$$

This proves that if δ is smaller than some critical value δ_1 and $\|\mathbf{y} - \mathbf{y}_0\| < \epsilon_\delta$, then the map $F_{\mathbf{y}}$ invariants any closed ball $K := \overline{B_\delta(\mathbf{x}_0)}$, i.e. $F_{\mathbf{y}}(K) \subset K$.

Now we want to show that choosing δ even smaller, the map $F_{\mathbf{y}}$ becomes a contraction. Indeed, from its definition in (6.5) we have:

$$F_{\mathbf{y}}(\mathbf{u}) - F_{\mathbf{y}}(\mathbf{w}) = -[D\mathbf{f}(\mathbf{x}_0)]^{-1}(\mathbf{g}(\mathbf{u}) - \mathbf{g}(\mathbf{w})),$$

or

$$\|F_{\mathbf{y}}(\mathbf{u}) - F_{\mathbf{y}}(\mathbf{w})\| \leq \|[D\mathbf{f}(\mathbf{x}_0)]^{-1}\|_{\text{HS}} \|\mathbf{g}(\mathbf{u}) - \mathbf{g}(\mathbf{w})\|, \quad \forall \mathbf{u}, \mathbf{w} \in \overline{B_{\delta}(\mathbf{x}_0)}.$$

Use again (6.4) with $\beta = \frac{1}{2(1+\|[D\mathbf{f}(\mathbf{x}_0)]^{-1}\|_{\text{HS}})}$: we obtain some $\delta_2 < \delta_1$ such that $F_{\mathbf{y}}(\overline{B_{\delta_2}(\mathbf{x}_0)}) \subset \overline{B_{\delta_2}(\mathbf{x}_0)}$ and

$$\|F_{\mathbf{y}}(\mathbf{u}) - F_{\mathbf{y}}(\mathbf{w})\| \leq \frac{1}{2}\|\mathbf{u} - \mathbf{w}\|, \quad \forall \mathbf{u}, \mathbf{w} \in \overline{B_{\delta_2}(\mathbf{x}_0)}.$$

Banach's fixed point theorem states that there exists a unique solution $\mathbf{x}_y \in \overline{B_{\delta_2}(\mathbf{x}_0)}$ which solves the equation $F_{\mathbf{y}}(\mathbf{x}) = \mathbf{x}$. Going back to (6.9) we conclude that if $\|\mathbf{y} - \mathbf{f}(\mathbf{x}_0)\| < \epsilon_{\delta_2}$, then there exists a solution $\mathbf{x}_y \in \overline{B_{\delta_2}(\mathbf{x}_0)}$ which solves the equation $\mathbf{f}(\mathbf{x}_y) = \mathbf{y}$. Since \mathbf{f} is injective on its domain, this solution is also unique. Moreover, $B_{\epsilon_{\delta_2}}(\mathbf{y}_0) \subset V$. Since \mathbf{y}_0 was arbitrary, V is open.

Let us now prove (ii). For any $\mathbf{y} \in V$ we constructed $\mathbf{x}_y = \mathbf{f}^{-1}(\mathbf{y})$ which solves $\mathbf{f}(\mathbf{x}_y) = \mathbf{y}$ and $F_{\mathbf{y}}(\mathbf{x}_y) = \mathbf{x}_y$. The fixed point equation rewrites as:

$$\mathbf{x}_y - \mathbf{x}_0 = [D\mathbf{f}(\mathbf{x}_0)]^{-1}(\mathbf{y} - \mathbf{y}_0) - [D\mathbf{f}(\mathbf{x}_0)]^{-1}(\mathbf{g}(\mathbf{x}_y) - \mathbf{g}(\mathbf{x}_0)). \quad (6.10)$$

We know that as soon as $\|\mathbf{y} - \mathbf{y}_0\| < \epsilon_{\delta_2}$, the point \mathbf{x}_y belongs to the ball around \mathbf{x}_0 where $\|\mathbf{g}(\mathbf{x}_y) - \mathbf{g}(\mathbf{x}_0)\| \leq \frac{1}{2(1+\|[D\mathbf{f}(\mathbf{x}_0)]^{-1}\|_{\text{HS}})}\|\mathbf{x}_y - \mathbf{x}_0\|$. Using this in (6.10) we get:

$$\|\mathbf{x}_y - \mathbf{x}_0\| \leq \|[D\mathbf{f}(\mathbf{x}_0)]^{-1}\|_{\text{HS}} \|\mathbf{y} - \mathbf{y}_0\| + \frac{1}{2}\|\mathbf{x}_y - \mathbf{x}_0\|,$$

or $\|\mathbf{x}_y - \mathbf{x}_0\| \leq 2\|[D\mathbf{f}(\mathbf{x}_0)]^{-1}\|_{\text{HS}} \|\mathbf{y} - \mathbf{y}_0\|$ for $\|\mathbf{y} - \mathbf{y}_0\|$ smaller than some critical value ϵ_{δ_2} . In other words, it means that

$$\lim_{\mathbf{y} \rightarrow \mathbf{y}_0} \mathbf{f}^{-1}(\mathbf{y}) = \mathbf{f}^{-1}(\mathbf{y}_0), \quad \|\mathbf{f}^{-1}(\mathbf{y}) - \mathbf{f}^{-1}(\mathbf{y}_0)\| \leq C\|\mathbf{y} - \mathbf{y}_0\|,$$

which shows that \mathbf{f}^{-1} is continuous on V . Moreover, (6.4) and the above estimate show that

$$\lim_{\mathbf{y} \rightarrow \mathbf{y}_0} \frac{\|\mathbf{g}(\mathbf{x}_y) - \mathbf{g}(\mathbf{x}_0)\|}{\|\mathbf{y} - \mathbf{y}_0\|} = 0.$$

Finally, we conclude from (6.10) that

$$\lim_{\mathbf{y} \rightarrow \mathbf{y}_0} \frac{\|\mathbf{f}^{-1}(\mathbf{y}) - \mathbf{f}^{-1}(\mathbf{y}_0) - [D\mathbf{f}(\mathbf{x}_0)]^{-1}(\mathbf{y} - \mathbf{y}_0)\|}{\|\mathbf{y} - \mathbf{y}_0\|} = 0$$

which shows that $[D\mathbf{f}^{-1}(\mathbf{y}_0)] = [D\mathbf{f}(\mathbf{x}_0)]^{-1}$ for every pair $\mathbf{f}(\mathbf{x}_0) = \mathbf{y}_0$, and we are done. \square

7 The implicit function theorem

In this section $d = m + n$ with $1 \leq m, n < d$. A vector $\mathbf{x} \in \mathbb{R}^d$ can be uniquely decomposed as $\mathbf{x} = [\mathbf{u}, \mathbf{w}]$ with $\mathbf{u} \in \mathbb{R}^m$ and $\mathbf{w} \in \mathbb{R}^n$.

Lemma 7.1. *Assume that D is a $d \times d$ matrix which has the following triangular form:*

$$D = \begin{bmatrix} A & B \\ 0_{n \times m} & I_{n \times n} \end{bmatrix},$$

where A is an $m \times m$ matrix and B is an arbitrary $m \times n$ matrix. Then D is invertible if and only if A is invertible and

$$D^{-1} = \begin{bmatrix} A^{-1} & -A^{-1}B \\ 0_{n \times m} & I_{n \times n} \end{bmatrix}. \quad (7.1)$$

Proof. We check by direct computation that $DD^{-1} = D^{-1}D = I_{d \times d}$. \square

Lemma 7.2. *Let $U \subset \mathbb{R}^d$ be an open set and $\mathbf{h} : U \mapsto \mathbb{R}^m$ be a $C^1(U; \mathbb{R}^m)$ function. Denote by $[D_{\mathbf{u}}\mathbf{h}](\mathbf{x})$ the partial $m \times m$ Jacobi matrix when \mathbf{w} is kept fixed, and by $[D_{\mathbf{w}}\mathbf{h}](\mathbf{x})$ the partial $m \times n$ Jacobi matrix when \mathbf{u} is kept fixed. Define the function $\mathbf{f} : U \mapsto \mathbb{R}^d$ by the formula $\mathbf{f}([\mathbf{u}, \mathbf{w}]) = [\mathbf{h}([\mathbf{u}, \mathbf{w}]), \mathbf{w}]$. Then $\mathbf{f} \in C^1(U; \mathbb{R}^d)$ and*

$$[D\mathbf{f}](\mathbf{x}) = \begin{bmatrix} [D_{\mathbf{u}}\mathbf{h}](\mathbf{x}) & [D_{\mathbf{w}}\mathbf{h}](\mathbf{x}) \\ 0_{n \times m} & I_{n \times n} \end{bmatrix}. \quad (7.2)$$

Proof. Direct computation. \square

We can now formulate the implicit function theorem.

Theorem 7.3. *Let $U \subset \mathbb{R}^d$ be an open set and $\mathbf{h} : U \mapsto \mathbb{R}^m$ be a $C^1(U; \mathbb{R}^m)$ function. Assume that there exists a point $\mathbf{a} = [\mathbf{u}_{\mathbf{a}}, \mathbf{w}_{\mathbf{a}}] \in U$ such that $\mathbf{h}(\mathbf{a}) = 0$ and the $m \times m$ partial Jacobi matrix $[D_{\mathbf{u}}\mathbf{h}](\mathbf{a})$ is invertible. Then there exists an open set $E \subset \mathbb{R}^n$ containing $\mathbf{w}_{\mathbf{a}}$, and a map $\mathbf{g} : E \mapsto \mathbb{R}^m$ in $C^1(E; \mathbb{R}^m)$, such that $\mathbf{g}(\mathbf{w}_{\mathbf{a}}) = \mathbf{u}_{\mathbf{a}}$ and $\mathbf{h}([\mathbf{g}(\mathbf{w}), \mathbf{w}]) = 0$ for all $\mathbf{w} \in E$. Moreover, $[D_{\mathbf{u}}\mathbf{h}]([\mathbf{g}(\mathbf{w}), \mathbf{w}])$ is invertible for all $\mathbf{w} \in E$ and*

$$[D\mathbf{g}](\mathbf{w}) = -[D_{\mathbf{u}}\mathbf{h}]^{-1}([\mathbf{g}(\mathbf{w}), \mathbf{w}]) [D_{\mathbf{w}}\mathbf{h}](([\mathbf{g}(\mathbf{w}), \mathbf{w}])). \quad (7.3)$$

Proof. As in Lemma 7.2 we define $\mathbf{f}([\mathbf{u}, \mathbf{w}]) = [\mathbf{h}([\mathbf{u}, \mathbf{w}]), \mathbf{w}]$ on U . From Lemma 7.1 and (7.2) we conclude that $[D\mathbf{f}](\mathbf{a})$ is invertible. The inverse function theorem 6.7 provides us with a ball $B_r(\mathbf{a})$ where \mathbf{f} is injective, $[D\mathbf{f}](\mathbf{x})$ is invertible on $B_r(\mathbf{a})$, and the set $V = \mathbf{f}(B_r(\mathbf{a}))$ is open in \mathbb{R}^d . Of course, the vector $[0, \mathbf{w}_{\mathbf{a}}] = \mathbf{f}(\mathbf{a}) \in V$. Since V is open, there exists $\epsilon > 0$ such that the d dimensional ball $B_\epsilon(\mathbf{f}(\mathbf{a})) = \{\mathbf{y} \in \mathbb{R}^d : \|\mathbf{y} - [0, \mathbf{w}_{\mathbf{a}}]\| < \epsilon\} \subset V$.

In general, if $[\mathbf{u}, \mathbf{w}] \in V$, assume that $\mathbf{f}^{-1}([\mathbf{u}, \mathbf{w}]) = [\mathbf{u}', \mathbf{w}'] \in B_r(\mathbf{a})$. Then we must have:

$$[\mathbf{u}, \mathbf{w}] = \mathbf{f}(\mathbf{f}^{-1}([\mathbf{u}, \mathbf{w}])) = [\mathbf{h}([\mathbf{u}', \mathbf{w}']), \mathbf{w}']$$

which shows that $\mathbf{w} = \mathbf{w}'$ and $\mathbf{u} = \mathbf{h}([\mathbf{u}', \mathbf{w}])$. This proves that $\mathbf{f}^{-1}([\mathbf{u}, \mathbf{w}])$ is also of the form $[\mathbf{u}', \mathbf{w}]$.

Let us define $\tilde{E} := \{\mathbf{w} \in \mathbb{R}^n : [0, \mathbf{w}] \in B_\epsilon(\mathbf{f}(\mathbf{a}))\}$, which is nothing but the n dimensional open ball $B_\epsilon(\mathbf{w}_{\mathbf{a}}) \subset \mathbb{R}^n$. Then for every $\mathbf{w} \in \tilde{E}$ we have that $[0, \mathbf{w}] \in V$ and

$$[0, \mathbf{w}] = \mathbf{f}(\mathbf{f}^{-1}([0, \mathbf{w}])) = [\mathbf{h}(\mathbf{f}^{-1}([0, \mathbf{w}])), \mathbf{w}]. \quad (7.4)$$

We already know that $\mathbf{f}^{-1}([0, \mathbf{w}])$ must be a vector of the form $[\mathbf{u}', \mathbf{w}]$, where \mathbf{u}' is nothing but the vector obtained from the first m components of $\mathbf{f}^{-1}([0, \mathbf{w}])$. Denote it by $\mathbf{g}(\mathbf{w})$. Then (7.4) implies that $\mathbf{h}([\mathbf{g}(\mathbf{w}), \mathbf{w}]) = 0$ on \tilde{E} . Moreover,

$$[\mathbf{g}(\mathbf{w}_{\mathbf{a}}), \mathbf{w}_{\mathbf{a}}] = \mathbf{f}^{-1}([0, \mathbf{w}_{\mathbf{a}}]) = \mathbf{a} = [\mathbf{u}_{\mathbf{a}}, \mathbf{w}_{\mathbf{a}}]$$

which implies $\mathbf{g}(\mathbf{w}_{\mathbf{a}}) = \mathbf{u}_{\mathbf{a}}$. Finally, since $(\mathbf{g}(\mathbf{w}))_j = (\mathbf{f}^{-1}([0, \mathbf{w}]))_j$ for $1 \leq j \leq m$ and $\mathbf{f}^{-1}([0, \mathbf{w}])$ is a $C^1(\tilde{E}; \mathbb{R}^d)$ map, then \mathbf{g} is $C^1(\tilde{E}; \mathbb{R}^m)$. Moreover, the determinant $\det[D_{\mathbf{u}}\mathbf{h}](\mathbf{g}(\mathbf{w}), \mathbf{w})$ is continuous on \tilde{E} and $\det[D_{\mathbf{u}}\mathbf{h}](\mathbf{a}) \neq 0$ because $[D_{\mathbf{u}}\mathbf{h}](\mathbf{a})$ is invertible. Thus we can find an open ball $E \subset \tilde{E}$ containing the point \mathbf{a} where the determinant $\det[D_{\mathbf{u}}\mathbf{h}](\mathbf{g}(\mathbf{w}), \mathbf{w})$ is different from zero, which implies that $[D_{\mathbf{u}}\mathbf{h}](\mathbf{g}(\mathbf{w}), \mathbf{w})$ is invertible on E . Finally, the formula (7.3) can be easily obtained by applying the chain rule to the equality $\mathbf{h}(\mathbf{g}(\mathbf{w}), \mathbf{w}) = 0$. \square

8 Brouwer's fixed point theorem

We say that $K \subset \mathbb{R}^d$ is convex if for every $\mathbf{x}, \mathbf{y} \in K$ we have that $(1-t)\mathbf{x} + t\mathbf{y} \in K$ for all $0 \leq t \leq 1$. A set K is called a convex body if K is convex, compact, and with at least one interior point.

Theorem 8.1. *Let $K \subset \mathbb{R}^d$ be a convex body. Let $\mathbf{f} : K \mapsto K$ be a continuous function which invariates K . Then \mathbf{f} has a (not necessarily unique) fixed point, that is a point $\mathbf{x} \in K$ such that $\mathbf{f}(\mathbf{x}) = \mathbf{x}$.*

Proof. The first thing we do is to reduce the problem from a general convex body to the unit ball in \mathbb{R}^d . We will show that there exists a bijection $\varphi : K \mapsto \overline{B_1(0)}$, which is continuous and with continuous inverse (a homeomorphism). If this is true, then it is enough to show that the function $\varphi \circ \mathbf{f} \circ \varphi^{-1} : \overline{B_1(0)} \mapsto \overline{B_1(0)}$ has a fixed point $\mathbf{a} \in \overline{B_1(0)}$. In that case, $\mathbf{x} = \varphi^{-1}(\mathbf{a}) \in K$.

Lemma 8.2. *Any convex body in \mathbb{R}^d is homeomorphic with the closed unit ball $\overline{B_1(0)}$.*

Proof. Let \mathbf{x}_0 be an interior point of K . There exists $r > 0$ such that $\overline{B_r(\mathbf{x}_0)} \subset K$. Define the continuous map $g : K \mapsto \mathbb{R}^d$ given by $g(\mathbf{x}) := (\mathbf{x} - \mathbf{x}_0)r^{-1}$. Define $\tilde{K} := g(K)$. It is easy to see that \tilde{K} is a convex and compact set. Moreover, the function $g : K \mapsto \tilde{K}$ is invertible and $g^{-1}(\mathbf{y}) = r\mathbf{y} + \mathbf{x}_0$. Both g and g^{-1} are continuous, and for every $\mathbf{y} \in \mathbb{R}^d$ with $\|\mathbf{y}\| \leq 1$ we have that $r\mathbf{y} + \mathbf{x}_0 \in \overline{B_r(\mathbf{x}_0)} \subset K$, thus $\mathbf{y} \in \tilde{K}$. This shows that $\overline{B_1(0)} \subset \tilde{K}$, thus \tilde{K} is a convex body containing the closed unit ball.

We now construct a homeomorphism between \tilde{K} and $\overline{B_1(0)}$. The boundary of \tilde{K} is denoted by $\partial\tilde{K}$, equals $\tilde{K} \setminus \text{int}(\tilde{K})$, and is a closed and bounded set. The boundary of $\overline{B_1(0)}$ is denoted by S^{d-1} and equals the unit sphere in \mathbb{R}^d . The ray connecting a given $\mathbf{x} \in \partial\tilde{K}$ with the origin intersects S^{d-1} in a point; in this way we define the function $h : \partial\tilde{K} \mapsto S^{d-1}$ where $h(\mathbf{x})$ is given by the above intersection.

The function h is injective; let us show this. Given $\mathbf{x} \in \partial\tilde{K}$, the cone $C(\mathbf{x})$ defined by joining \mathbf{x} with all the points of $\overline{B_1(0)}$ must belong to the convex set \tilde{K} . But $C(\mathbf{x})$ also contains the ray joining \mathbf{x} with 0, and all the points of the segment strictly between \mathbf{x} and 0 are interior points of $C(\mathbf{x})$, thus interior points of \tilde{K} . It means that no two different points of $\partial\tilde{K}$ can be placed on the same ray starting from the origin, which proves the injectivity of h .

Let us show that the function h is also surjective. Consider any ray generated by $\hat{x} \in S^{d-1}$, starting from the origin and parametrized by $R(\hat{x}) := \{\lambda\hat{x} : \lambda \geq 0\}$. Consider the set $E(\hat{x}) := R(\hat{x}) \cap \text{int}(\tilde{K})$. This set must be bounded, because \tilde{K} is bounded. Thus the set of non-negative real numbers $\{\|\mathbf{y}\| : \mathbf{y} \in E(\hat{x})\} \subset \mathbb{R}$ is bounded, thus it has a supremum $c < \infty$. The supremum is an accumulation point, thus there must exist a sequence of points $\{\mathbf{y}_n\}_{n \geq 1} \subset E(\hat{x})$ such that $\|\mathbf{y}_n\| \rightarrow c$. But this sequence is also contained by the compact set $\tilde{K} \cap R(\hat{x})$. It means that there exists a subsequence \mathbf{y}_{n_k} which converges to some point $\mathbf{u} \in \tilde{K} \cap R(\hat{x})$, i.e. $\|\mathbf{y}_{n_k} - \mathbf{u}\| \rightarrow 0$ when $k \rightarrow \infty$. Thus we also have $\|\mathbf{y}_{n_k}\| \rightarrow \|\mathbf{u}\|$ which shows that $\|\mathbf{u}\| = c$. Now \mathbf{u} cannot be an interior point of \tilde{K} , because in that case we could find points of $E(\hat{x})$ which are farther away from the origin than \mathbf{u} , contradicting the maximality of $\|\mathbf{u}\|$. Thus $\mathbf{u} \in \partial\tilde{K}$, which proves that the ray hits at least one point of the boundary. Hence h is surjective.

Thus h is bijective and invertible. The continuity of h is obvious.

Let us now prove that $h^{-1} : S^{d-1} \mapsto \partial\tilde{K}$ is also continuous. For every $\hat{x} \in S^{d-1}$, the point $h^{-1}(\hat{x})$ is the unique point of $\partial\tilde{K}$ which is hit by the ray defined by \hat{x} . Assume that h^{-1} is not continuous at some $\hat{a} \in S^{d-1}$. It means that we can find some $\epsilon_0 > 0$ and a sequence $\{\hat{x}_n\}_{n \geq 1} \subset S^{d-1}$, such that $\hat{x}_n \rightarrow \hat{a}$ and $\|h^{-1}(\hat{x}_n) - h^{-1}(\hat{a})\| \geq \epsilon_0$. The vector $h^{-1}(\hat{x}_n)$ is parallel with \hat{x}_n and the same is true for the pair $h^{-1}(\hat{a})$ and \hat{a} . Thus if n is large enough, the last inequality implies that either $\|h^{-1}(\hat{x}_n)\| \leq \|h^{-1}(\hat{a})\| - \epsilon_0/2$ or $\|h^{-1}(\hat{a})\| + \epsilon_0/2 \leq \|h^{-1}(\hat{x}_n)\|$. Assume that there are infinitely many cases where the first situation holds true. Then if n is large enough, the point $h^{-1}(\hat{x}_n)$ enters in the cone $C(h^{-1}(\hat{a}))$ and must be an interior point of \tilde{K} , contradiction. In the other situation, $h^{-1}(\hat{a})$ would eventually become an interior element of the cone $C(h^{-1}(\hat{x}_n))$ for large enough n , again contradiction.

Let us define the map $\phi : \tilde{K} \mapsto \overline{B_1(0)}$ by $\phi(\mathbf{x}) := \mathbf{x}/\|h^{-1}(\mathbf{x}/\|\mathbf{x}\|)\|$ if $\mathbf{x} \neq 0$ and $\phi(0) = 0$. It is nothing but taking \mathbf{x} and dividing it with the length of the segment between 0 and the point on

the boundary corresponding to the ray generated by $\mathbf{x}/\|\mathbf{x}\|$. Clearly, ϕ is continuous. It is easy to check that the inverse of ϕ is given by $\phi^{-1} : \overline{B_1(0)} \mapsto K$ where $\phi^{-1}(\mathbf{y}) := \mathbf{y}\|h^{-1}(\mathbf{y}/\|\mathbf{y}\|)\|$ if $\mathbf{y} \neq 0$ and $\phi^{-1}(0) = 0$. This inverse is also continuous since h is.

In conclusion, $\varphi := \phi \circ g : K \mapsto \overline{B_1(0)}$ is a homeomorphism, and we are done. \square

Thus from now on we will assume without loss of generality that $K = \overline{B_1(0)}$. And also from now we will assume that there exists a continuous function which invariates $B_1(0)$ and has no fixed points.

Lemma 8.3. *Assume that $\mathbf{f} : \overline{B_1(0)} \mapsto \overline{B_1(0)}$ is continuous with no fixed points. Then there exists a smooth function $\tilde{f} : \overline{B_1(0)} \mapsto \overline{B_1(0)}$ with the same property.*

Proof. Our assumption says that $\|\mathbf{f}(\mathbf{x}) - \mathbf{x}\| > 0$ for all $\mathbf{x} \in \overline{B_1(0)}$. The real valued map

$$\overline{B_1(0)} \ni \mathbf{x} \mapsto \|\mathbf{f}(\mathbf{x}) - \mathbf{x}\| \in \mathbb{R}$$

is continuous and defined on a compact set. Thus it attains its minimum in some point \mathbf{x}_m . It follows that:

$$\|\mathbf{f}(\mathbf{x}) - \mathbf{x}\| \geq \|\mathbf{f}(\mathbf{x}_m) - \mathbf{x}_m\| = \epsilon_0 > 0. \quad (8.1)$$

Let us extend \mathbf{f} to the whole of \mathbb{R}^d in the following way. Define $g : \mathbb{R}^d \mapsto \overline{B_1(0)}$ by $g(\mathbf{x}) = \mathbf{f}(\mathbf{x})$ if $\|\mathbf{x}\| \leq 1$, and $g(\mathbf{x}) = \mathbf{f}(\mathbf{x}/\|\mathbf{x}\|)$ if $\|\mathbf{x}\| > 1$. Clearly the extension is continuous, and $\|g(\mathbf{x})\| \leq 1$ for all \mathbf{x} .

Define the function $j : \mathbb{R}^d \mapsto \mathbb{R}$, $j(\mathbf{x}) = e^{-1/(1-\|\mathbf{x}\|^2)}$ if $\|\mathbf{x}\| < 1$ and $j(\mathbf{x}) = 0$ if $\|\mathbf{x}\| \geq 1$. The function j is non-negative, belongs to $C^\infty(\mathbb{R}^d)$ and has a positive integral $I := \int_{\mathbb{R}^d} j(\mathbf{x}) d\mathbf{x} > 0$. Define $\tilde{j}(\mathbf{x}) := j(\mathbf{x})/I$. Then $\int_{\mathbb{R}^d} \tilde{j}(\mathbf{x}) d\mathbf{x} = 1$.

Now if $\delta > 0$ we define the function $J_\delta(\mathbf{x}) := \delta^{-d} \tilde{j}(\delta^{-1}\mathbf{x})$. Clearly, J_δ is non-negative, belongs to $C^\infty(\mathbb{R}^d)$, it is non-zero only if $\|\mathbf{x}\| < \delta$, and $\int_{\mathbb{R}^d} J_\delta(\mathbf{x}) d\mathbf{x} = 1$ independently of δ .

Define the function $g_\delta : \overline{B_1(0)} \mapsto \overline{B_1(0)}$ by the formula:

$$g_\delta(\mathbf{x}) := \int_{\mathbb{R}^d} J_\delta(\mathbf{x} - \mathbf{y}) g(\mathbf{y}) d\mathbf{y} = \int_{\mathbb{R}^d} g(\mathbf{x} - \mathbf{y}) J_\delta(\mathbf{y}) d\mathbf{y}. \quad (8.2)$$

That the range of g_δ is included in $\overline{B_1(0)}$ is a consequence of the fact that $\|g(\mathbf{y})\| \leq 1$ and $\int_{\mathbb{R}^d} J_\delta(\mathbf{x} - \mathbf{y}) d\mathbf{y} = 1$ independently of \mathbf{x} . The function g_δ is smooth because J_δ is smooth.

Now we can write:

$$g_\delta(\mathbf{x}) - g(\mathbf{x}) = \int_{\mathbb{R}^d} [g(\mathbf{x} - \mathbf{y}) - g(\mathbf{x})] J_\delta(\mathbf{y}) d\mathbf{y} = \int_{\|\mathbf{y}\| \leq \delta} [g(\mathbf{x} - \mathbf{y}) - g(\mathbf{x})] J_\delta(\mathbf{y}) d\mathbf{y}, \quad (8.3)$$

where the second equality comes from the support properties of J_δ . If we impose the condition $\delta < 1$, then $\mathbf{x} - \mathbf{y} \in \overline{B_2(0)}$ if $\|\mathbf{y}\| \leq \delta$ and $\|\mathbf{x}\| \leq 1$. The function g restricted to the compact set $\overline{B_2(0)}$ is uniformly continuous, thus there exists some $\delta_0 > 0$ small enough such that

$$\|g(\mathbf{x}') - g(\mathbf{x}'')\| \leq \epsilon_0/2 \quad \text{whenever} \quad \|\mathbf{x}' - \mathbf{x}''\| \leq \delta_0, \quad \mathbf{x}', \mathbf{x}'' \in \overline{B_2(0)}.$$

Applying this estimate in (8.3) we obtain that $\|g_{\delta_0}(\mathbf{x}) - g(\mathbf{x})\| \leq \epsilon_0/2$, for all $\mathbf{x} \in \overline{B_1(0)}$. Using this in (8.1) it follows:

$$\|g_{\delta_0}(\mathbf{x}) - \mathbf{x}\| \geq \epsilon_0/2 > 0, \quad \forall \mathbf{x} \in \overline{B_1(0)}. \quad (8.4)$$

The function g_{δ_0} is our \tilde{f} and the proof of this lemma is over. \square

From now on we can assume that our function \mathbf{f} is smooth and with no fixed points in $\overline{B_1(0)}$. The next lemma shows that such a function \mathbf{f} would allow us to construct a smooth retraction of the unit ball onto its boundary.

Lemma 8.4. *Assume that $\mathbf{f} : \overline{B_1(0)} \mapsto \overline{B_1(0)}$ is smooth with no fixed points. Then there exists a smooth function $\mathbf{h} : \overline{B_1(0)} \mapsto S^{d-1}$ such that $\mathbf{h}(\mathbf{x}) = \mathbf{x}$ if $\mathbf{x} \in S^{d-1}$.*

Proof. We know that there exists some $\epsilon_0 > 0$ such that $\|\mathbf{f}(\mathbf{x}) - \mathbf{x}\| \geq \epsilon_0$ for all $\mathbf{x} \in \overline{B_1(0)}$. We define the unit vector $\mathbf{w}(\mathbf{x}) := (\|\mathbf{x} - \mathbf{f}(\mathbf{x})\|)^{-1} (\mathbf{x} - \mathbf{f}(\mathbf{x}))$ which defines the direction of a straight line starting in $\mathbf{f}(\mathbf{x})$ and going through \mathbf{x} . This line is parametrized as $\mathbf{f}(\mathbf{x}) + t\mathbf{w}(\mathbf{x})$, with $t \geq 0$. The value $t = \|\mathbf{x} - \mathbf{f}(\mathbf{x})\|$ gives \mathbf{x} . For even larger values of t we approach the boundary. There exists a unique positive value of $t(\mathbf{x}) \geq \|\mathbf{x} - \mathbf{f}(\mathbf{x})\| \geq \epsilon_0$ which corresponds to the intersection of this line with the unit sphere S^{d-1} . Namely, from the condition $\|\mathbf{f}(\mathbf{x}) + t\mathbf{w}(\mathbf{x})\|^2 = 1$ we obtain:

$$t(\mathbf{x}) = -\mathbf{f}(\mathbf{x}) \cdot \mathbf{w}(\mathbf{x}) + \sqrt{(\mathbf{f}(\mathbf{x}) \cdot \mathbf{w}(\mathbf{x}))^2 + 1 - \|\mathbf{f}(\mathbf{x})\|^2} \geq \|\mathbf{x} - \mathbf{f}(\mathbf{x})\|,$$

where $\mathbf{f}(\mathbf{x}) \cdot \mathbf{w}(\mathbf{x})$ is the inner product in \mathbb{R}^d . The only problem related to the smoothness of this function could appear if the square root can be zero. The square root is zero if $\|\mathbf{f}(\mathbf{x})\| = 1$ and $0 = \mathbf{f}(\mathbf{x}) \cdot \mathbf{w}(\mathbf{x})$. Equivalently, $\mathbf{f}(\mathbf{x}) \cdot \mathbf{x} = 1$. The last equality demands that $\mathbf{x} = \mathbf{f}(\mathbf{x})$, both vectors having unit length and sitting on the boundary, situation excluded by our assumption of absence of fixed points. Thus $t(\mathbf{x})$ is smooth, and we can define

$$\mathbf{h}(\mathbf{x}) := \mathbf{f}(\mathbf{x}) + t(\mathbf{x}) \mathbf{w}(\mathbf{x}) \in S^{d-1}$$

which ends the proof. □

Lemma 8.5. *Assume that $\mathbf{h} : \overline{B_1(0)} \mapsto S^{d-1}$ is smooth and $\mathbf{h}(\mathbf{x}) = \mathbf{x}$ if $\mathbf{x} \in S^{d-1}$. If $0 \leq s \leq 1$, define the map $\mathbf{g}_s : \overline{B_1(0)} \mapsto \overline{B_1(0)}$ given by $\mathbf{g}_s(\mathbf{x}) = (1-s)\mathbf{x} + s\mathbf{h}(\mathbf{x})$. Then there exists $0 < s_0 < 1$ such that \mathbf{g}_s is a bijection for all $0 \leq s \leq s_0$.*

Proof. First of all, we note that if $\mathbf{x} \in S^{d-1}$ then $\mathbf{g}_s(\mathbf{x}) = \mathbf{x}$. Thus the only thing we need to show is that \mathbf{g}_s is injective and $\mathbf{g}_s(B_1(0)) = B_1(0)$.

For the injectivity part: consider the equality $\mathbf{g}_s(\mathbf{x}) = \mathbf{g}_s(\mathbf{y})$ for some $\mathbf{x}, \mathbf{y} \in \overline{B_1(0)}$. This can be rewritten as:

$$\mathbf{x} - \mathbf{y} = -\frac{s}{1-s}(\mathbf{h}(\mathbf{x}) - \mathbf{h}(\mathbf{y})).$$

Reasoning as in Lemma 6.3 we can find a constant $C_h > 0$ such that $\|\mathbf{h}(\mathbf{u}) - \mathbf{h}(\mathbf{w})\| \leq C_h \|\mathbf{u} - \mathbf{w}\|$ for all $\mathbf{u}, \mathbf{w} \in \overline{B_1(0)}$. Thus we obtain:

$$\|\mathbf{x} - \mathbf{y}\| \leq \frac{C_h s}{1-s} \|\mathbf{x} - \mathbf{y}\|$$

which imposes $\mathbf{x} = \mathbf{y}$ if s is smaller than some small enough value $0 < \tilde{s} < 1$.

Now let us assume that $0 \leq s \leq \tilde{s}$. We want to prove that there exists $0 < s_0 \leq \tilde{s}$ such that $\mathbf{g}_s(B_1(0)) = B_1(0)$ for all $0 \leq s \leq s_0$.

One inclusion is easy: if $\|\mathbf{x}\| < 1$, then $\|\mathbf{g}_s(\mathbf{x})\| \leq (1-s)\|\mathbf{x}\| + s < 1$. Thus $\mathbf{g}_s(B_1(0)) \subset B_1(0)$.

The other inclusion is more complicated. Let us consider the equation $\mathbf{g}_s(\mathbf{x}) = \mathbf{z}$, where $\|\mathbf{z}\| \leq 1/4$ is arbitrary. This equation can be rewritten as $\mathbf{x} = (1-s)^{-1}\{\mathbf{z} - s\mathbf{h}(\mathbf{x})\}$. Now if s is smaller than some small enough value s_1 , the vector $T(\mathbf{x}) := (1-s)^{-1}\mathbf{z} - s(1-s)^{-1}\mathbf{h}(\mathbf{x})$ obeys $\|T(\mathbf{x})\| \leq 1/2$ for all $\|\mathbf{x}\| \leq 1$. In particular, T invariates $\overline{B_{\frac{1}{2}}(0)}$. Also:

$$\|T(\mathbf{u}) - T(\mathbf{w})\| \leq C_h s \|\mathbf{u} - \mathbf{w}\|, \quad \forall \mathbf{u}, \mathbf{w} \in \overline{B_{\frac{1}{2}}(0)}.$$

Thus if $s < s_2 := \min\{s_1, C_h^{-1}\}$, the map T is a contraction and has a unique fixed point. This fixed point solves the equation $\mathbf{g}_s(\mathbf{x}) = \mathbf{z}$. Thus until now we showed that

$$\overline{B_{\frac{1}{4}}(0)} \subset \mathbf{g}_s(B_1(0)), \quad 0 \leq s < s_2 < 1.$$

Another important observation which we have to prove is that $\mathbf{g}_s(B_1(0))$ is an open set. Indeed, we have $[D\mathbf{g}_s(\mathbf{x})] = (1-s)I_{d \times d} + s[Dh(\mathbf{x})]$ and $\det[D\mathbf{g}_s(\mathbf{x})] \geq 1/2$ if s is smaller than some small enough s_3 , for all $\mathbf{x} \in B_1(0)$; let $\mathbf{y} = \mathbf{g}_s(\mathbf{a})$ for some $\mathbf{a} \in B_1(0)$. Then from Theorem 6.7 (i) it follows that there is some r small enough such that $\mathbf{g}_s(B_r(\mathbf{a}))$ is open, and since $\mathbf{y} \in \mathbf{g}_s(B_r(\mathbf{a}))$ there exists $\epsilon > 0$ so that $B_\epsilon(\mathbf{g}_s(\mathbf{a})) \subset \mathbf{g}_s(B_r(\mathbf{a})) \subset \mathbf{g}_s(B_1(0))$.

Now fix $0 < s_0 < \min\{s_2, s_3\}$. For $0 \leq s \leq s_0$ we know that $\mathbf{g}_s(B_1(0))$ is open and $\overline{B_{\frac{1}{4}}(0)} \subset \mathbf{g}_s(B_1(0)) \subset B_1(0)$. We need to show that $B_1(0) \subset \mathbf{g}_s(B_1(0))$.

Assume the contrary: there exists some $\mathbf{y}_0 \in B_1(0)$ which does not belong to $\mathbf{g}_s(B_1(0))$. Denote by I the closed segment joining 0 with \mathbf{y}_0 . The set $E := I \cap \mathbf{g}_s(B_1(0))$ is not empty. Moreover, the set:

$$\{\|\mathbf{y}\| : \mathbf{y} \in I \cap \mathbf{g}_s(B_1(0))\} \subset [0, \|\mathbf{y}_0\|]$$

is not empty, and has a supremum $c < 1$. The supremum is always a limit point, hence there exists a sequence $\{\mathbf{y}_n\}_{n \geq 1} \subset I \cap \mathbf{g}_s(B_1(0))$ such that $\|\mathbf{y}_n\| \rightarrow c$. Because I is compact, there exists a subsequence \mathbf{y}_{n_k} which converges in I to some point $\tilde{\mathbf{y}} \in I$, thus $\tilde{\mathbf{y}}$ is an adherent point of $\mathbf{g}_s(B_1(0))$ and $\|\tilde{\mathbf{y}}\| = c < 1$. Clearly, $\tilde{\mathbf{y}} \notin \mathbf{g}_s(B_1(0))$ because otherwise, since $\mathbf{g}_s(B_1(0))$ is open, we could extend $I \cap \mathbf{g}_s(B_1(0))$ even further away from the origin, contradicting the maximality of the length of $\tilde{\mathbf{y}}$.

Thus we have constructed $\tilde{\mathbf{y}} \in \overline{\mathbf{g}_s(B_1(0))} \setminus \mathbf{g}_s(B_1(0))$ with $\|\tilde{\mathbf{y}}\| \leq \|\mathbf{y}_0\| < 1$. Being an adherent point of $\mathbf{g}_s(B_1(0))$, there must exist a sequence $\{\mathbf{z}_n\}_{n \geq 1} \subset \mathbf{g}_s(B_1(0))$ such that $\mathbf{z}_n \rightarrow \tilde{\mathbf{y}}$. There exists a sequence $\{\mathbf{x}_n\}_{n \geq 1} \subset B_1(0)$ such that $\mathbf{g}_s(\mathbf{x}_n) = \mathbf{z}_n$. We can find a subsequence \mathbf{x}_{n_k} which converges to some $\mathbf{x}_0 \in \overline{B_1(0)}$. Since $\mathbf{g}_s(\mathbf{x}_{n_k}) = \mathbf{z}_{n_k} \rightarrow \tilde{\mathbf{y}}$ and due to the continuity of \mathbf{g}_s , we must have $\mathbf{g}_s(\mathbf{x}_0) = \tilde{\mathbf{y}}$. But since $\tilde{\mathbf{y}} \notin \mathbf{g}_s(B_1(0))$, it must be that $\mathbf{x}_0 \in S^{d-1}$. But on the boundary, $\mathbf{g}_s(\mathbf{x}_0) = \mathbf{x}_0$, which contradicts our assumption that $\|\tilde{\mathbf{y}}\| \leq \|\mathbf{y}_0\| < 1$. Therefore, \mathbf{y}_0 cannot exist, and $B_1(0) \subset \mathbf{g}_s(B_1(0))$. □

We are finally ready to prove Brouwer's theorem. In the previous lemma we considered the smooth map $\mathbf{g}_s : \overline{B_1(0)} \mapsto \overline{B_1(0)}$. Define the function:

$$F(s) := \int_{B_1(0)} \det[D\mathbf{g}_s(\mathbf{x})] d\mathbf{x}, \quad 0 \leq s \leq 1.$$

The determinant of the Jacobi matrix $[D\mathbf{g}_s(\mathbf{x})]$ is a polynomial in s , thus $F(s)$ is a polynomial. Moreover, we have shown that if $0 \leq s \leq s_0$, the map \mathbf{g}_s is nothing but a smooth and bijective change of coordinates in $B_1(0)$ with $\det[D\mathbf{g}_s(\mathbf{x})] > 0$, thus $F(s)$ is constant on $[0, s_0]$ and equal to the volume of $B_1(0)$. But if a polynomial is locally constant, then it is constant everywhere. Thus $F(1)$ should also be equal to the volume of $B_1(0)$.

Now let us show that this cannot be true. If $s = 1$, then $\mathbf{g}_1(\mathbf{x}) = \mathbf{h}(\mathbf{x})$ on $B_1(0)$. It means that

$$1 = \|\mathbf{h}(\mathbf{x})\|^2 = \mathbf{g}_1(\mathbf{x}) \cdot \mathbf{g}_1(\mathbf{x}) = \sum_{k=1}^d (\mathbf{g}_1(\mathbf{x}))_k^2.$$

Differentiating with respect to x_j we obtain

$$0 = \sum_{k=1}^d [\partial_j (\mathbf{g}_1(\mathbf{x}))_k] (\mathbf{g}_1(\mathbf{x}))_k, \quad 1 \leq j \leq d,$$

or $[D\mathbf{g}_1(\mathbf{x})]^* \mathbf{g}_1(\mathbf{x}) = 0$ for all \mathbf{x} . Since $\|\mathbf{g}_1(\mathbf{x})\| = 1$, we have that $[D\mathbf{g}_1(\mathbf{x})]^*$ is not injective, thus not invertible, hence with zero determinant. Therefore $\det[D\mathbf{g}_1(\mathbf{x})] = \det[D\mathbf{g}_1(\mathbf{x})]^* = 0$ for all \mathbf{x} , and $F(1) = 0 \neq \text{vol}(B_1(0))$. This contradiction can be traced back to our assumption which claimed that \mathbf{f} has no fixed points. The proof is over. □

9 Schauder's fixed point theorem

Theorem 9.1. *Let X be a Banach space, and let $K \subset X$ be a non-empty, compact, and convex set. Then given any continuous mapping $f: K \mapsto K$ there exists $x \in K$ such that $f(x) = x$.*

Proof. Given $\epsilon > 0$, the family of open sets $\{B_\epsilon(x) : x \in K\}$ is an open covering of K . Because K is compact, there exists a finite subcover, i.e. there exists N points p_1, \dots, p_N of K such that the balls $B_\epsilon(p_i)$ cover the whole set K .

Let K_ϵ be the convex hull of p_1, \dots, p_N , defined by:

$$K_\epsilon := \left\{ \sum_{j=1}^N t_j p_j, \quad \sum_{j=1}^N t_j = 1, \quad t_j \geq 0 \right\} \subset K.$$

It is an easy computation to show that K_ϵ is a convex set. Moreover, K_ϵ is a finite dimensional object, immersed in an at most $N-1$ dimensional Euclidian space generated by the vectors $p_j - p_1$, where $j \in \{2, 3, \dots, N\}$.

Define the function $g_j : K \mapsto \mathbb{R}_+$ by $g_j(x) = \epsilon - \|x - p_j\|$ if $x \in B_\epsilon(p_j)$, and $g_j(x) = 0$ otherwise. Each function g_j is continuous, while $g(x) = \sum_{j=1}^N g_j(x)$ is positive due to the fact that any x has to be in some ball, where the corresponding g_j is positive. Since g is continuous and K compact, there exists $\delta > 0$ such that $g(x) \geq \delta$ for every $x \in K$.

Now consider the continuous map $\pi_\epsilon : K \rightarrow K_\epsilon$ given by:

$$\pi_\epsilon(x) := \sum_{j=1}^N \frac{g_j(x)}{g(x)} p_j, \quad \sum_{j=1}^N \frac{g_j(x)}{g(x)} = 1.$$

Since $\|g_j(x)(x - p_j)\| \leq g_j(x)\epsilon$ for all j , we have:

$$\|\pi_\epsilon(x) - x\| \leq \sum_{j=1}^N \frac{\|g_j(x)(p_j - x)\|}{g(x)} \leq \epsilon, \quad \forall x \in K. \quad (9.1)$$

Now we define:

$$f_\epsilon : K_\epsilon \rightarrow K_\epsilon, \quad f_\epsilon(x) = \pi_\epsilon(f(x)).$$

This is a continuous function defined on a convex and compact set K_ϵ in a finite dimensional vector space. By Brouwer's fixed point theorem it admits a fixed point x_ϵ

$$f_\epsilon(x_\epsilon) = x_\epsilon.$$

Using (9.1) we get:

$$\|\pi_\epsilon(f(x_\epsilon)) - f(x_\epsilon)\| \leq \epsilon,$$

thus for every $\epsilon > 0$ we have constructed $x_\epsilon \in K_\epsilon \subset K$ such that $\|f(x_\epsilon) - x_\epsilon\| \leq \epsilon$.

Choosing $1/n$ instead of ϵ , we construct a sequence $\{x_n\}_{n \geq 1} \subset K$ such that $\|f(x_n) - x_n\| \leq 1/n$. Since K is sequentially compact, we can find a subsequence x_{n_k} which converges to some point $\bar{x} \in K$ when $k \rightarrow \infty$. By writing:

$$\|f(\bar{x}) - \bar{x}\| \leq \|f(\bar{x}) - f(x_{n_k})\| + \|f(x_{n_k}) - x_{n_k}\| + \|x_{n_k} - \bar{x}\|, \quad k \geq 1,$$

we observe that due to the continuity of f at \bar{x} , the right hand side tends to zero with k . Thus $f(\bar{x}) = \bar{x}$ and we are done. \square

10 Kakutani's fixed point theorem

Let $A \subset \mathbb{R}^d$ be a closed set, and denote by 2^A the set of all subsets of A . We say that $F : A \mapsto 2^A$ is upper semicontinuous if the following property holds: assume that $\{\mathbf{x}_n\}_{n \geq 1} \subset A$ with $\mathbf{x}_n \rightarrow \mathbf{x}_\infty \in A$, $\{\mathbf{y}_n\}_{n \geq 1} \subset A$ with $\mathbf{y}_n \rightarrow \mathbf{y}_\infty \in A$, and $\mathbf{y}_n \in F(\mathbf{x}_n)$; then we must have $\mathbf{y}_\infty \in F(\mathbf{x}_\infty)$.

Note that if we choose $\mathbf{x}_n = \mathbf{x}_\infty$ for all n , then the upper semicontinuity implies that if $\mathbf{y}_n \rightarrow \mathbf{y}_\infty \in A$ and $\mathbf{y}_n \in F(\mathbf{x}_\infty)$, then $\mathbf{y}_\infty \in F(\mathbf{x}_\infty)$. In other words, $F(\mathbf{x}_\infty)$ is always closed for all $\mathbf{x}_\infty \in A$.

Theorem 10.1. *Let $K \subset \mathbb{R}^d$ be a convex body. Let $F : K \mapsto 2^K$ be upper semicontinuous, such that $F(\mathbf{x}) \subset K$ is convex and nonempty. Then there exists $\mathbf{x}^* \in K$ such that $\mathbf{x}^* \in F(\mathbf{x}^*)$.*

Proof. Since K is compact, for every $m \geq 1$ there exist N_m points denoted by $\{\mathbf{w}_{j,m}\}_{j=1}^{N_m}$ such that $K \subset \bigcup_{j=1}^{N_m} B_{\frac{1}{m}}(\mathbf{w}_{j,m})$. It is important for what follows to note that we may choose the points $\mathbf{w}_{j,m}$ such that each ball $B_{\frac{1}{m}}(\mathbf{x})$ contains at most \mathcal{N} (only depending on the dimension d) points $\mathbf{w}_{j,m}$, independent of m and \mathbf{x} .

For every $1 \leq j \leq N_m$ we define a map $g_{j,m} : K \mapsto \mathbb{R}_+$ by $g_{j,m}(\mathbf{x}) = \epsilon - \|\mathbf{x} - \mathbf{w}_{j,m}\|$ if $\mathbf{x} \in B_{\frac{1}{m}}(\mathbf{w}_{j,m})$, and $g_{j,m}(\mathbf{x}) = 0$ otherwise. Each function $g_{j,m}$ is continuous, while $g_m(\mathbf{x}) = \sum_{j=1}^{N_m} g_{j,m}(\mathbf{x})$ is positive due to the fact that any \mathbf{x} has to be in some ball j , where the corresponding $g_{j,m}$ is positive. Since g_m is continuous and K compact, there exists $\delta > 0$ such that $g_m(\mathbf{x}) \geq \delta$ for every $\mathbf{x} \in K$.

For every $1 \leq j \leq N_m$ we choose some $\mathbf{y}_{j,m} \in F(\mathbf{w}_{j,m}) \subset K$ in an arbitrary way. Define the map

$$\mathbf{f}_m : K \mapsto K, \quad \mathbf{f}_m(\mathbf{x}) := \sum_{j=1}^{N_m} \frac{g_{j,m}(\mathbf{x})}{g_m(\mathbf{x})} \mathbf{y}_{j,m}, \quad \sum_{j=1}^{N_m} \frac{g_{j,m}(\mathbf{x})}{g_m(\mathbf{x})} = 1, \quad \mathbf{y}_j \in F(\mathbf{w}_{j,m}).$$

The function \mathbf{f}_m is continuous and defined on a convex body, thus Brouwer's fixed point theorem provides us with a fixed point $\mathbf{x}_m \in K$ such that $\mathbf{f}_m(\mathbf{x}_m) = \sum_{j=1}^{N_m} \frac{g_{j,m}(\mathbf{x}_m)}{g_m(\mathbf{x}_m)} \mathbf{y}_{j,m} = \mathbf{x}_m$, for every $m \geq 1$.

Because K is sequentially compact, we may find a subsequence \mathbf{x}_{m_k} which converges to some $\mathbf{x}^* \in K$ when $k \rightarrow \infty$. It is important to note that when j varies from 1 to N_{m_k} we have $g_{j,m_k}(\mathbf{x}_{m_k}) \neq 0$ only for those \mathbf{w}_{j,m_k} which obey $\|\mathbf{x}_{m_k} - \mathbf{w}_{j,m_k}\| \leq \frac{1}{m_k} \leq \frac{1}{k}$. There are at most \mathcal{N} indexes j such that $g_{j,m_k}(\mathbf{x}_{m_k}) \neq 0$:

$$\mathbf{f}_{m_k}(\mathbf{x}_{m_k}) = \mathbf{x}_{m_k} = \sum_{\|\mathbf{x}_{m_k} - \mathbf{w}_{j,m_k}\| \leq \frac{1}{m_k}} \frac{g_{j,m_k}(\mathbf{x}_{m_k})}{g_{m_k}(\mathbf{x}_{m_k})} \mathbf{y}_{j,m_k}.$$

Thus for a fixed $k \geq 1$ we have finitely many \mathbf{w}_{j,m_k} (at most \mathcal{N} , independently of k) which all lie in a small ball of radius $1/m_k$ around \mathbf{x}_{m_k} . We can reorganize the \mathcal{N} closest points \mathbf{w}_{j,m_k} and their corresponding \mathbf{y}_{j,m_k} as \mathcal{N} pairs of sequences $\{\tilde{\mathbf{w}}_k^s\}_{k \geq 1}$ and $\tilde{\mathbf{y}}_k^s$, with $\lim_{k \rightarrow \infty} \|\tilde{\mathbf{w}}_k^s - \mathbf{x}^*\| = 0$ and $\tilde{\mathbf{y}}_k^s \in F(\tilde{\mathbf{w}}_k^s)$ for all $1 \leq s \leq \mathcal{N}$. With the new notation:

$$\mathbf{f}_{m_k}(\mathbf{x}_{m_k}) = \mathbf{x}_{m_k} = \sum_{s=1}^{\mathcal{N}} \frac{g_{s,m_k}(\mathbf{x}_{m_k})}{g_{m_k}(\mathbf{x}_{m_k})} \tilde{\mathbf{y}}_k^s, \quad \lim_{k \rightarrow \infty} \tilde{\mathbf{w}}_k^s = \mathbf{x}^*.$$

Now we can choose a subsequence $\tilde{\mathbf{y}}_{k_n}^s$, $n \geq 1$, such that $\tilde{\mathbf{y}}_{k_n}^s$ converges to some $\mathbf{y}^s \in K$. Thus:

$$\mathbf{x}_{m_{k_n}} = \sum_{s=1}^{\mathcal{N}} \frac{g_{s,m_{k_n}}(\mathbf{x}_{m_{k_n}})}{g_{m_{k_n}}(\mathbf{x}_{m_{k_n}})} \tilde{\mathbf{y}}_{k_n}^s, \quad \lim_{n \rightarrow \infty} \tilde{\mathbf{w}}_{k_n}^s = \mathbf{x}^*, \quad \lim_{n \rightarrow \infty} \tilde{\mathbf{y}}_{k_n}^s = \mathbf{y}^s.$$

To summarize, we have $\lim_{n \rightarrow \infty} \tilde{\mathbf{w}}_{k_n}^s = \mathbf{x}^*$, $\lim_{n \rightarrow \infty} \tilde{\mathbf{y}}_{k_n}^s = \mathbf{y}^s$, and $\tilde{\mathbf{y}}_{k_n}^s \in F(\tilde{\mathbf{w}}_{k_n}^s)$. Because F is upper semicontinuous, we must have $\mathbf{y}^s \in F(\mathbf{x}^*)$ for all $1 \leq s \leq \mathcal{N}$. Moreover,

$$\left\| \mathbf{x}_{m_{k_n}} - \sum_{s=1}^{\mathcal{N}} \frac{g_{s,m_{k_n}}(\mathbf{x}_{m_{k_n}})}{g_{m_{k_n}}(\mathbf{x}_{m_{k_n}})} \mathbf{y}^s \right\| \leq \max_{s=1}^{\mathcal{N}} \|\tilde{\mathbf{y}}_{k_n}^s - \mathbf{y}^s\| \rightarrow 0.$$

Since $F(\mathbf{x}^*)$ is convex, and all $\mathbf{y}^s \in F(\mathbf{x}^*)$, the convex combination $\sum_{s=1}^{\mathcal{N}} \frac{g_{s, m_{k_n}}(\mathbf{x}_{m_{k_n}})}{g_{m_{k_n}}(\mathbf{x}_{m_{k_n}})} \mathbf{y}^s$ is an element of $F(\mathbf{x}^*)$. Since $\mathbf{x}_{m_{k_n}} \rightarrow \mathbf{x}^*$, we have that \mathbf{x}^* must be an adherent point of $F(\mathbf{x}^*)$. Since $F(\mathbf{x}^*)$ is closed, then $\mathbf{x}^* \in F(\mathbf{x}^*)$ and we are done. \square

11 Existence of Nash equilibrium for finite games with two players

In order to simplify notation, we only consider two players. Let us start with a few definitions. Let d_1 and d_2 be two natural numbers larger than 1. Consider two finite sets $S_i = \{s_{1,i}, s_{2,i}, \dots, s_{d_i,i}\}$, $i \in \{1, 2\}$, where the element $s_{k,i}$ is called the k 'th pure strategy of player i . The sets S_1 and S_2 are called the sets of pure strategies.

A payoff function of player i is just an arbitrary non-negative function $\pi^i : S_1 \times S_2 \mapsto \mathbb{R}_+$. It is completely characterized by the non-negative numbers $\pi^{i,jk} := \pi^i(s_{1,j}, s_{2,k})$ which define a $d_1 \times d_2$ matrix. The set of mixed strategies for player i , denoted by M_i , is the set of all probability distributions defined on S_i . We can write:

$$M_i := \{\rho_i : S_i \mapsto \mathbb{R}_+ : \sum_{k=1}^{d_i} \rho_i(s_{k,i}) = 1\}.$$

It is easy to see that each such probability distribution is completely characterized by its d_i values taken on the different possible pure strategies. Hence we have the identification with a compact set (bounded and closed) in \mathbb{R}^{d_i} :

$$\tilde{M}_i := \{\mathbf{x} \in \mathbb{R}^{d_i} : 0 \leq x_k \leq 1, \sum_{k=1}^{d_i} x_k = 1\}, \quad i \in \{1, 2\}.$$

The expected payoff to player i is defined to be the function:

$$K := \tilde{M}_1 \times \tilde{M}_2 \ni \mathbf{z} = [\mathbf{x}, \mathbf{y}] \mapsto v_i(\mathbf{z}) := \sum_{j=1}^{d_1} \sum_{k=1}^{d_2} x_j y_k \pi^{i,jk} \in \mathbb{R}_+.$$

A point $\mathbf{z}_e = [\mathbf{x}_e, \mathbf{y}_e] \in K \subset \mathbb{R}^{d_1+d_2}$ is called a Nash equilibrium if $v_1(\mathbf{z}_e) \geq v_1([\mathbf{x}, \mathbf{y}_e])$ and $v_2(\mathbf{z}_e) \geq v_2([\mathbf{x}_e, \mathbf{y}])$ for every $\mathbf{x} \in \tilde{M}_1$ and $\mathbf{y} \in \tilde{M}_2$.

Theorem 11.1. *Every finite game has a Nash equilibrium.*

Proof. We will use the Kakutani fixed point theorem. For every $\mathbf{z} = [\mathbf{x}, \mathbf{y}] \in K$ we define the 'best response functions':

$$K \ni [x, y] \mapsto \phi_1(\mathbf{z}) := \{\tilde{\mathbf{x}} \in \tilde{M}_1 : v_1([\tilde{\mathbf{x}}, \mathbf{y}]) \geq v_1([\mathbf{w}, \mathbf{y}]), \quad \forall \mathbf{w} \in \tilde{M}_1\} \in 2^{\tilde{M}_1}$$

and

$$K \ni [x, y] \mapsto \phi_2(\mathbf{z}) := \{\tilde{\mathbf{y}} \in \tilde{M}_2 : v_2([\mathbf{x}, \tilde{\mathbf{y}}]) \geq v_2([\mathbf{x}, \mathbf{u}]), \quad \forall \mathbf{u} \in \tilde{M}_2\} \in 2^{\tilde{M}_2},$$

and $F : K \mapsto 2^K$ given by $F(\mathbf{z}) := \phi_1(\mathbf{z}) \times \phi_2(\mathbf{z})$.

Now let us assume that F has a fixed point in the sense of Kakutani's theorem. It means that there exists $\mathbf{z}_e \in K$ such that $\mathbf{z}_e \in F(\mathbf{z}_e)$. In other words, $\mathbf{x}_e \in \phi_1(\mathbf{z}_e)$ which is the same as $v_1([\mathbf{x}_e, \mathbf{y}_e]) \geq v_1([\mathbf{w}, \mathbf{y}_e]), \forall \mathbf{w} \in \tilde{M}_1$, and $\mathbf{y}_e \in \phi_2(\mathbf{z}_e)$ which is the same as $v_2([\mathbf{x}_e, \mathbf{y}_e]) \geq v_2([\mathbf{x}_e, \mathbf{u}]), \forall \mathbf{u} \in \tilde{M}_2$. These are exactly the Nash equilibrium conditions.

The rest of the proof will show that F obeys the conditions of Kakutani's theorem. First, $K \subset \mathbb{R}^{d_1+d_2}$ is (sequentially) compact because it is the cartesian product of two sequentially compact sets.

Second, let us show that $F(z)$ is convex, with $z = [\mathbf{x}, \mathbf{y}]$. Let $[\mathbf{u}_1, \mathbf{w}_1]$ and $[\mathbf{u}_2, \mathbf{w}_2]$ in $F(\mathbf{z})$. For every $0 \leq t \leq 1$ we have to show that $(1-t)\mathbf{u}_1 + t\mathbf{u}_2 \in \phi_1(\mathbf{z})$ and $(1-t)\mathbf{w}_1 + t\mathbf{w}_2 \in \phi_2(\mathbf{z})$. We only show the first property. Since \mathbf{u}_1 and \mathbf{u}_2 belong to $\phi_1(\mathbf{z})$, according to the definition we have:

$$v_1([\mathbf{u}_1, \mathbf{y}]) \geq v_1([\mathbf{w}, \mathbf{y}]) \text{ and } v_1([\mathbf{u}_2, \mathbf{y}]) \geq v_1([\mathbf{w}, \mathbf{y}]), \quad \forall \mathbf{w} \in \tilde{M}_1.$$

Since v_1 is linear in \mathbf{u} for fixed \mathbf{y} , we have:

$$v_1([(1-t)\mathbf{u}_1 + t\mathbf{u}_2, \mathbf{y}]) = (1-t)v_1([\mathbf{u}_1, \mathbf{y}]) + tv_1([\mathbf{u}_2, \mathbf{y}]) \geq v_1([\mathbf{w}, \mathbf{y}]), \quad \forall \mathbf{w} \in \tilde{M}_1$$

which shows that $(1-t)\mathbf{u}_1 + t\mathbf{u}_2 \in \phi_1(\mathbf{z})$.

Third, we need to show that F is upper semicontinuous. Let $\mathbf{z}_n \rightarrow \mathbf{z}_\infty \in K$, $\mathbf{f}_n \rightarrow \mathbf{f}_\infty \in K$, and $\mathbf{f}_n \in F(\mathbf{z}_n)$ for all $n \geq 1$. We need to show that $\mathbf{f}_\infty \in F(\mathbf{f}_\infty)$. But this is an easy consequence of the fact that v_1 and v_2 are continuous functions which preserve inequalities at the limit. The proof is over. \square

12 The Hairy Ball Theorem

The question we want to answer here is of geometric nature: given the unit sphere $S^{d-1} := \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\| = 1\} \subset \mathbb{R}^d$ for $d \geq 2$, is it possible to find a continuous tangent vector field $\tilde{\mathbf{w}} : S^{d-1} \mapsto \mathbb{R}^d$ which vanishes nowhere? If it is possible, then $\|\tilde{\mathbf{w}}(\mathbf{x})\|$ has a positive lower bound because S^{d-1} is compact, and in this way we would be able to construct a continuous vector field $\mathbf{w} : S^{d-1} \mapsto S^{d-1}$ given by $\mathbf{w}(\mathbf{x}) := \frac{1}{\|\tilde{\mathbf{w}}(\mathbf{x})\|} \tilde{\mathbf{w}}(\mathbf{x})$, which satisfies $\mathbf{w}(\mathbf{x}) \cdot \mathbf{x} = 0$.

If $d = 2p$ is even, then such a vector field exists and let us construct an example. If $\mathbf{x} = [x_1, x_2, \dots, x_p, x_{p+1}, \dots, x_{2p-1}, x_{2p}]$ then we can define $\mathbf{u} \in \mathbb{R}^p$ by $\mathbf{u} := [x_1, x_2, \dots, x_p]$ and $\tilde{\mathbf{u}} \in \mathbb{R}^p$ by $\tilde{\mathbf{u}} := [x_{p+1}, \dots, x_{2p-1}, x_{2p}]$. Thus $\mathbf{x} = [\mathbf{u}, \tilde{\mathbf{u}}]$. Define $\mathbf{w}(\mathbf{x}) = [-\tilde{\mathbf{u}}, \mathbf{u}]$. Then clearly $\|\mathbf{w}(\mathbf{x})\| = \|\mathbf{x}\| = 1$ and $\mathbf{w}(\mathbf{x}) \cdot \mathbf{x} = 0$.

Theorem 12.1. *If $d \geq 3$ is odd, then one cannot construct a continuous map $\mathbf{w} : S^{d-1} \mapsto S^{d-1}$ such that $\mathbf{w}(\mathbf{x}) \cdot \mathbf{x} = 0$ on S^{d-1} .*

Proof. We will assume that such a vector field exists, and then we will arrive at a contradiction.

Lemma 12.2. *Let $d \geq 2$. Assume that $\mathbf{w} : S^{d-1} \mapsto S^{d-1}$ is continuous and $\mathbf{w}(\mathbf{x}) \cdot \mathbf{x} = 0$ for all $\mathbf{x} \in S^{d-1}$. Define $D_1 := \{\mathbf{x} \in \mathbb{R}^d : \frac{1}{2} < \|\mathbf{x}\| < \frac{3}{2}\}$. Then there exists a smooth map $\tilde{\mathbf{w}} : D_1 \mapsto \mathbb{R}^d$ such that $\tilde{\mathbf{w}}(D_1) \subset S^{d-1}$ and $\tilde{\mathbf{w}}(\mathbf{x}) \cdot \mathbf{x} = 0$ for all $\mathbf{x} \in D_1$.*

Proof. If $\mathbf{x} \neq 0$ we denote by $\hat{\mathbf{x}} := \frac{1}{\|\mathbf{x}\|} \mathbf{x} \in S^{d-1}$. Define $D_2 := \{\mathbf{x} \in \mathbb{R}^d : \frac{1}{4} \leq \|\mathbf{x}\| \leq 2\}$, and consider the function $\mathbf{w}_2 : D_2 \mapsto \mathbb{R}^d$ given by $\mathbf{w}_2(\mathbf{x}) := \mathbf{w}(\hat{\mathbf{x}})$. Then \mathbf{w}_2 is continuous on the compact set D_2 , thus uniformly continuous. Given any $\epsilon > 0$ we may find $\delta > 0$ such that $\|\mathbf{w}_2(\mathbf{x}) - \mathbf{w}_2(\mathbf{x}')\| < \epsilon$ as soon as $\mathbf{x}, \mathbf{x}' \in D_2$ and $\|\mathbf{x} - \mathbf{x}'\| < \delta$. We will reason as in Lemma 8.3: consider the function J_δ and define as in (8.2)

$$\mathbf{g}_\delta(\mathbf{x}) := \int_{\mathbb{R}^d} J_\delta(\mathbf{x} - \mathbf{y}) \mathbf{w}_2(\mathbf{y}) d\mathbf{y} = \int_{\mathbb{R}^d} \mathbf{w}_2(\mathbf{x} - \mathbf{y}) J_\delta(\mathbf{y}) d\mathbf{y}, \quad \mathbf{x} \in D_1.$$

As in that lemma, we may write

$$\mathbf{g}_\delta(\mathbf{x}) - \mathbf{w}_2(\mathbf{x}) = \int_{\mathbb{R}^d} [\mathbf{w}_2(\mathbf{x} - \mathbf{y}) - \mathbf{w}_2(\mathbf{x})] J_\delta(\mathbf{y}) d\mathbf{y} = \int_{\|\mathbf{y}\| \leq \delta} [\mathbf{w}_2(\mathbf{x} - \mathbf{y}) - \mathbf{w}_2(\mathbf{x})] J_\delta(\mathbf{y}) d\mathbf{y}.$$

If $\delta < 1/4$ then $\mathbf{x} - \mathbf{y} \in D_2$, hence using the uniform continuity of \mathbf{w}_2 on D_2 we may find $\delta_0 < 1/4$ small enough such that $\|\mathbf{w}_2(\mathbf{x} - \mathbf{y}) - \mathbf{w}_2(\mathbf{x})\| \leq 1/10$ for every $\mathbf{x} \in D_1$ and $\|\mathbf{y}\| \leq \delta_0$. This leads to

$$\|\mathbf{g}_{\delta_0}(\mathbf{x}) - \mathbf{w}_2(\mathbf{x})\| \leq 1/10 \quad \text{and} \quad \|\mathbf{g}_{\delta_0}(\mathbf{x})\| \geq 9/10, \quad \forall \mathbf{x} \in D_1.$$

The function $\mathbf{w}_3 : D_1 \mapsto \mathbb{R}^d$ given by (remember that $\mathbf{w}_2(\mathbf{x}) \cdot \mathbf{x} = 0$)

$$\mathbf{w}_3(\mathbf{x}) := \mathbf{g}_{\delta_0}(\mathbf{x}) - \mathbf{x} \frac{\mathbf{g}_{\delta_0}(\mathbf{x}) \cdot \mathbf{x}}{\|\mathbf{x}\|^2} = \mathbf{g}_{\delta_0}(\mathbf{x}) - \mathbf{x} \frac{[\mathbf{g}_{\delta_0}(\mathbf{x}) - \mathbf{w}_2(\mathbf{x})] \cdot \mathbf{x}}{\|\mathbf{x}\|^2}$$

is smooth on D_1 and obeys:

$$\mathbf{w}_3(\mathbf{x}) \cdot \mathbf{x} = 0, \quad \|\mathbf{w}_3(\mathbf{x})\| \geq \|\mathbf{g}_{\delta_0}(\mathbf{x})\| - \|\mathbf{g}_{\delta_0}(\mathbf{x}) - \mathbf{w}_2(\mathbf{x})\| \geq 8/10, \quad \forall \mathbf{x} \in D_1.$$

Finally, we can define the function $\tilde{\mathbf{w}}(\mathbf{x}) := \frac{1}{\|\mathbf{w}_3(\mathbf{x})\|} \mathbf{w}_3(\mathbf{x}) \in S^{d-1}$ which is smooth and orthogonal on \mathbf{x} . Moreover, we have the estimate (see Lemma 6.3 for the notation):

$$\|\Delta \tilde{\mathbf{w}}\|_{\infty, \bar{D}_1} < \infty. \quad (12.1)$$

□

Define $D_3 := \{\mathbf{x} \in \mathbb{R}^d : \frac{9}{10} < \|\mathbf{x}\| < \frac{10}{9}\} \subset D_1$. If $s \in \mathbb{R}$ we denote by

$$E_s := \left\{ \mathbf{x} \in \mathbb{R}^d : \sqrt{\frac{81}{100} + s^2} < \|\mathbf{x}\| < \sqrt{\frac{100}{81} + s^2} \right\}.$$

Lemma 12.3. *Let $\mathbf{h}_s : D_3 \mapsto E_s$ given by $\mathbf{h}_s(\mathbf{x}) := \mathbf{x} + s\tilde{\mathbf{w}}(\mathbf{x})$. If $|s| > 0$ is sufficiently small, then the map \mathbf{h}_s is a bijection.*

Proof. Because $\|\mathbf{h}_s(\mathbf{x})\|^2 = \|\mathbf{x}\|^2 + s^2$ it is easy to see that $\mathbf{h}_s(D_3) \subset E_s$ for all s . We need to show that \mathbf{h}_s is injective and surjective if s is small enough.

Let us start by showing injectivity. Assume that \mathbf{h}_s is not injective in a neighborhood of $s = 0$. Then there exists a sequence $\{s_n\}_{n \geq 1}$ which converges to 0 such that for every $n \neq 1$ there exist $\mathbf{x}_n \neq \mathbf{y}_n \in D_3$ such that $\mathbf{h}_{s_n}(\mathbf{x}_n) = \mathbf{h}_{s_n}(\mathbf{y}_n)$. This is equivalent with $\mathbf{x}_n - \mathbf{y}_n = s_n[\tilde{\mathbf{w}}(\mathbf{y}_n) - \tilde{\mathbf{w}}(\mathbf{x}_n)]$, which implies that $\|\mathbf{x}_n - \mathbf{y}_n\| \leq 2|s_n|$. Since $D_3 \subset D_1$, if $|s_n|$ is small enough then the whole segment joining \mathbf{x}_n and \mathbf{y}_n is included in D_1 . Using (12.1) and (6.3) for $\tilde{\mathbf{w}}$ we get that

$$\|\mathbf{x}_n - \mathbf{y}_n\| = |s_n| \|\tilde{\mathbf{w}}(\mathbf{y}_n) - \tilde{\mathbf{w}}(\mathbf{x}_n)\| \leq |s_n| \|\Delta \tilde{\mathbf{w}}\|_{\infty, \bar{D}_1} \|\mathbf{x}_n - \mathbf{y}_n\|$$

which is incompatible with $\mathbf{x}_n \neq \mathbf{y}_n$ if $|s_n|$ is small enough.

Now we have to prove that \mathbf{h}_s is surjective if $|s|$ is small enough. Let $\mathbf{y} \in E_s$. We have to show that the equation $\mathbf{h}_s(\mathbf{x}) = \mathbf{y}$ has a solution. This equation is equivalent with $\mathbf{x} = \mathbf{y} - s\tilde{\mathbf{w}}(\mathbf{x})$, which looks like a fixed point equation.

Define the closed set $D_4 := \{\mathbf{x} \in \mathbb{R}^d : \frac{8}{9} \leq \|\mathbf{x}\| \leq \frac{9}{8}\}$. If $|s|$ is sufficiently small, then $E_s \subset D_4$. Moreover, $D_3 \subset D_4 \subset D_1$.

The set D_4 is closed in \mathbb{R}^d , thus together with the induced Euclidian metric it forms a complete metric space. We want to show that the map $\mathbf{f}_{\mathbf{y},s} : D_4 \mapsto D_4$ given by $\mathbf{f}_{\mathbf{y},s}(\mathbf{x}) := \mathbf{y} - s\tilde{\mathbf{w}}(\mathbf{x})$ is a contraction on D_4 provided $|s|$ is small enough. If this is true, then the unique fixed point which obeys $\mathbf{f}_{\mathbf{y},s}(\mathbf{x}) = \mathbf{x}$ has the property that $\|\mathbf{x}_y\|^2 + s^2 = \|\mathbf{y}\|^2$, thus $\mathbf{x}_y \in D_3$ if $\mathbf{y} \in E_s$.

Now let us show that $\mathbf{f}_{\mathbf{y},s}$ is a contraction for small $|s|$. First, using that $\|\tilde{\mathbf{w}}(\mathbf{x})\| = 1$, then if $\frac{10}{9} + 2|s| \leq \frac{9}{8}$ and $\frac{8}{9} \leq \frac{9}{10} - |s|$ we have that $\mathbf{f}_{\mathbf{y},s}(D_4) \subset D_4$. Second, if $\mathbf{x}_1, \mathbf{x}_2 \in D_4$ with $\|\mathbf{x}_1 - \mathbf{x}_2\| \geq \frac{1}{100}$, we have:

$$\|\mathbf{f}_{\mathbf{y},s}(\mathbf{x}_1) - \mathbf{f}_{\mathbf{y},s}(\mathbf{x}_2)\| \leq 200|s| \|\mathbf{x}_1 - \mathbf{x}_2\|, \quad \text{if } \|\mathbf{x}_1 - \mathbf{x}_2\| \geq \frac{1}{100}.$$

Third, if $\mathbf{x}_1, \mathbf{x}_2 \in D_4$ with $\|\mathbf{x}_1 - \mathbf{x}_2\| \leq \frac{1}{100}$, then since the ball $B_{\frac{1}{100}}(\mathbf{x}_1)$ is completely included in D_1 , the straight segment joining \mathbf{x}_1 and \mathbf{x}_2 is included in D_1 and using again (6.3) we obtain:

$$\|\mathbf{f}_{\mathbf{y},s}(\mathbf{x}_1) - \mathbf{f}_{\mathbf{y},s}(\mathbf{x}_2)\| \leq |s| \|\Delta \tilde{\mathbf{w}}\|_{\infty, \bar{D}_1} \|\mathbf{x}_1 - \mathbf{x}_2\|, \quad \text{if } \|\mathbf{x}_1 - \mathbf{x}_2\| \leq \frac{1}{100}.$$

Thus $\mathbf{f}_{\mathbf{y},s}$ is a contraction if $|s|$ is smaller than some critical value s_0 which is independent of \mathbf{y} . This implies that the fixed point exists for all $\mathbf{y} \in E_s$ provided $0 \leq |s| \leq s_0$, and we are done. □

We are now ready to finish the proof of the Hairy Ball Theorem. Since the map \mathbf{h}_s is a smooth bijection between D_3 and E_s and $\det[D\mathbf{h}_s](\mathbf{x}) > 0$ if $|s| \leq s_0$, we must have the equality:

$$\text{Vol}(E_s) = \int_{D_3} \det[D\mathbf{h}_s](\mathbf{x}) d\mathbf{x}, \quad |s| \leq s_0.$$

The right hand side of the above equality is a polynomial in s of degree at most d . The left hand side can be calculated explicitly: it equals the difference of the volumes of two d -dimensional balls:

$$\text{Vol}(E_s) = \frac{\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2} + 1)} \left\{ \left(\frac{100}{81} + s^2 \right)^{\frac{d}{2}} - \left(\frac{81}{100} + s^2 \right)^{\frac{d}{2}} \right\}.$$

This function is analytic around $s = 0$, and since it equals a polynomial if $|s| \leq s_0$, by analytic continuation it must be a polynomial (thus analytic) for all $s \in \mathbb{C}$. The function $\left(\frac{100}{81} + s^2\right)^{\frac{d}{2}}$ is analytic if $|s| < 10/9$, thus $\left(\frac{81}{100} + s^2\right)^{\frac{d}{2}}$ must also be analytic on the same disk. Since $d = 2p + 1$ is odd, the function $\left(\frac{81}{100} + s^2\right)^{\frac{d}{2}}$ can be factorized as the product $\left(\frac{81}{100} + s^2\right)^p \left(\frac{81}{100} + s^2\right)^{\frac{1}{2}}$. But this function is not analytic at $s = \frac{9i}{9}$, and this is our contradiction. \square

13 The Jordan Curve Theorem

13.1 Some preparatory results

Let (X, d) be a metric space with the topology generated by d . We say that X is not connected if we can find two non-empty open sets O_1 and O_2 which are disjoint and $X = O_1 \cup O_2$. A subset $A \subset X$ is not connected if the induced metric space (A, d) is not connected. If a metric space can be written as a union of connected sets $\{O_i\}$, then they are called the connected components of X .

Let $-\infty < a \leq b < \infty$ and let $\gamma : [a, b] \mapsto \mathbb{R}^2$ be a homeomorphism (continuous, invertible and with continuous inverse). Then $\gamma([a, b])$ is called an arc. A set $A \subset \mathbb{R}^2$ is called path-connected if given any two distinct points $\mathbf{x}, \mathbf{y} \in A$ we can find an arc joining them and which is included in A .

Lemma 13.1. *Let A be a non-empty connected open set in the Euclidian space \mathbb{R}^2 . Then A is path-connected.*

Proof. Choose a point $\mathbf{x} \in A$. Define the set $O_1 \subset A$ which contains all the points $\mathbf{y} \in A$ which can be connected with \mathbf{x} by an arc. Let us show that O_1 is open. If $\mathbf{y} \in O_1$ then there exists an arc $\gamma([a, b]) \subset A$ with $\gamma(a) = \mathbf{x}$ and $\gamma(b) = \mathbf{y}$. Since \mathbf{y} is an interior point of A , there exists $\epsilon > 0$ such that if $\|\mathbf{x}' - \mathbf{y}\| < \epsilon$ then $\mathbf{x}' \in A$. But all such points can be joined with \mathbf{y} by a straight line included in A . Thus \mathbf{x} can be joined with \mathbf{x}' by an arc included in A , hence $B_\epsilon(\mathbf{y}) \subset O_1$.

Now if $O_1 = A$ then we are done. If not, the set $O_2 := A \setminus O_1$ is not empty. One can prove in a similar manner that O_2 is open: if \mathbf{y} cannot be joined with \mathbf{x} by an arc included in A , then no points close enough to \mathbf{y} can be joined with \mathbf{x} . But since $A = O_1 \cup O_2$, it would mean that A is not connected, contradiction. \square

Lemma 13.2. *Let M be a compact set in the Euclidian space \mathbb{R}^2 . If $\mathbf{x} \in \mathbb{R}^2$ we define*

$$\mathbb{R}^2 \ni \mathbf{x} \mapsto d(\mathbf{x}, M) := \inf\{\|\mathbf{y} - \mathbf{x}\| : \mathbf{y} \in M\} \in \mathbb{R}.$$

Then this map is continuous.

Proof. For a fixed \mathbf{x} , the map $M \ni \mathbf{y} \mapsto \|\mathbf{y} - \mathbf{x}\| \in \mathbb{R}$ is continuous and defined on a compact set. Thus it attains its minimum at some point $\mathbf{y}_x \in M$. Hence for every $\mathbf{x} \in \mathbb{R}^2$ there exists $\mathbf{y}_x \in M$ such that $d(\mathbf{x}, M) = \|\mathbf{x} - \mathbf{y}_x\|$. We note the inequalities:

$$d(\mathbf{x}', M) \leq \|\mathbf{x}' - \mathbf{y}_x\| \leq \|\mathbf{x}' - \mathbf{x}\| + \|\mathbf{x} - \mathbf{y}_x\| = \|\mathbf{x}' - \mathbf{x}\| + d(\mathbf{x}, M),$$

which due to the symmetry lead to:

$$|d(\mathbf{x}, M) - d(\mathbf{x}', M)| \leq \|\mathbf{x} - \mathbf{x}'\|$$

for all $\mathbf{x}, \mathbf{x}' \in \mathbb{R}^2$, which proves Lipschitz continuity. \square

The next lemma is a poor man's version of the Tietze extension theorem. It roughly says that given an arc M in \mathbb{R}^2 contained in a large closed ball B , we can find a continuous function $g : B \mapsto M$ which extends the identity map on M .

Lemma 13.3. *Let M be an arc in \mathbb{R}^2 . Let $\mathbf{x}_0 \in \mathbb{R}^2$ and consider $\overline{B_r(\mathbf{x}_0)}$ a ball sufficiently large such that $M \subset \overline{B_r(\mathbf{x}_0)}$. Then there exists a continuous map $g : \overline{B_r(\mathbf{x}_0)} \mapsto M$ such that $g(\mathbf{x}) = \mathbf{x}$ if $\mathbf{x} \in M$.*

Proof. Since M is an arc, it is bounded and closed, thus compact. Moreover, M is homeomorphic with a closed interval in \mathbb{R} , thus we can find a continuous function $\gamma : [0, 1] \mapsto M$ with continuous inverse $\gamma^{-1} : M \mapsto [0, 1]$. If we can find a continuous function $F : \overline{B_r(\mathbf{x}_0)} \mapsto [0, 1]$ such that $F(\mathbf{x}) = \gamma^{-1}(\mathbf{x})$ for all $\mathbf{x} \in M$, then the extension we are looking for is $g = \gamma \circ F$.

Let us define the function F . If $\mathbf{x} \in M$ we put $F(\mathbf{x}) = \gamma^{-1}(\mathbf{x})$. If $\mathbf{x} \in \overline{B_r(\mathbf{x}_0)} \setminus M$ we put

$$F(\mathbf{x}) := \inf_{\mathbf{y} \in M} \left\{ \gamma^{-1}(\mathbf{y}) + \frac{\|\mathbf{x} - \mathbf{y}\|}{d(\mathbf{x}, M)} - 1 \right\}.$$

If $\mathbf{x} \notin M$, the map

$$M \ni \mathbf{y} \mapsto \gamma^{-1}(\mathbf{y}) + \frac{\|\mathbf{x} - \mathbf{y}\|}{d(\mathbf{x}, M)} - 1 \in \mathbb{R}$$

is continuous and defined on a compact set, thus there exists some $\mathbf{w}(\mathbf{x}) \in M$ such that

$$F(\mathbf{x}) = \gamma^{-1}(\mathbf{w}(\mathbf{x})) + \frac{\|\mathbf{x} - \mathbf{w}(\mathbf{x})\|}{d(\mathbf{x}, M)} - 1. \quad (13.2)$$

Let us show that the range of F is the interval $[0, 1]$. If $\mathbf{x} \in M$ it is obvious. If $\mathbf{x} \notin M$ then we know from Lemma 13.2 that there exists some $\mathbf{y}_x \in M$ such that $0 < \|\mathbf{x} - \mathbf{y}_x\| = d(\mathbf{x}, M) \leq \|\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{y} \in M$. Thus $0 \leq \gamma^{-1}(\mathbf{y}) + \frac{\|\mathbf{x} - \mathbf{y}\|}{d(\mathbf{x}, M)} - 1$, for all $\mathbf{y} \in M$, which implies that $0 \leq F(\mathbf{x})$. Moreover,

$$F(\mathbf{x}) \leq \gamma^{-1}(\mathbf{y}_x) + \frac{\|\mathbf{x} - \mathbf{y}_x\|}{d(\mathbf{x}, M)} - 1 = \gamma^{-1}(\mathbf{y}_x) \leq 1.$$

Now we want to prove that F is continuous. Let $\mathbf{a} \in M$. According to the definition of F , we have that $F(\mathbf{a}) = \gamma^{-1}(\mathbf{a})$. Consider any sequence $\{\mathbf{x}_n\} \subset \overline{B_r(\mathbf{x}_0)}$ which converges to \mathbf{a} . We can split it in a subsequence included in M , denoted by $\{\mathbf{x}_n^M\}$, and a subsequence in $\overline{B_r(\mathbf{x}_0)} \setminus M$, denoted by $\{\mathbf{x}_n^{M^c}\}$. Since $F(\mathbf{x}_n^M) = \gamma^{-1}(\mathbf{x}_n^M)$ and because γ^{-1} is continuous on M , we have $F(\mathbf{x}_n^M) \rightarrow F(\mathbf{a}) = \gamma^{-1}(\mathbf{a})$. What we have to prove now is that $F(\mathbf{x}_n^{M^c}) \rightarrow \gamma^{-1}(\mathbf{a})$ provided $\mathbf{x}_n^{M^c} \rightarrow \mathbf{a}$. We note that the sequence of minimizing points $\mathbf{w}(\mathbf{x}_n^{M^c})$ must converge to \mathbf{a} ; otherwise, since $d(\mathbf{x}_n^{M^c}, M) \leq \|\mathbf{x}_n^{M^c} - \mathbf{a}\| \rightarrow 0$, we would eventually have $F(\mathbf{x}_n^{M^c}) > 1$ for some subsequence. Moreover, any sequence $\mathbf{y}_{\mathbf{x}_n^{M^c}}$ defined by $d(\mathbf{x}_n^{M^c}, M) = \|\mathbf{x}_n^{M^c} - \mathbf{y}_{\mathbf{x}_n^{M^c}}\| \leq \|\mathbf{x}_n^{M^c} - \mathbf{a}\|$ must converge to \mathbf{a} . Then we have:

$$\gamma^{-1}(\mathbf{w}(\mathbf{x}_n^{M^c})) \leq F(\mathbf{x}_n^{M^c}) \leq \gamma^{-1}(\mathbf{y}_{\mathbf{x}_n^{M^c}}),$$

where the first inequality is a consequence of (13.2) and of $\|\mathbf{x}_n^{M^c} - \mathbf{w}(\mathbf{x}_n^{M^c})\| \geq d(\mathbf{x}_n^{M^c}, M)$, while the second inequality is a consequence of the definition of F . Then the continuity of γ^{-1} at \mathbf{a} ensures that $F(\mathbf{x}_n^M) \rightarrow \gamma^{-1}(\mathbf{a})$ and we are done.

Now let $\mathbf{a} \in \overline{B_r(\mathbf{x}_0)} \setminus M$ and consider any sequence $\mathbf{x}_n \rightarrow \mathbf{a}$. Since M is closed, we may consider that $\mathbf{x}_n \notin M$ for all n . Then we can write:

$$F(\mathbf{a}) = \gamma^{-1}(\mathbf{w}(\mathbf{a})) + \frac{\|\mathbf{a} - \mathbf{w}(\mathbf{a})\|}{d(\mathbf{a}, M)} - 1, \quad F(\mathbf{x}_n) = \gamma^{-1}(\mathbf{w}(\mathbf{x}_n)) + \frac{\|\mathbf{x}_n - \mathbf{w}(\mathbf{x}_n)\|}{d(\mathbf{x}_n, M)} - 1.$$

From the definition of F and using the triangle inequality we have:

$$\begin{aligned} F(\mathbf{a}) &\leq \gamma^{-1}(\mathbf{w}(\mathbf{x}_n)) + \frac{\|\mathbf{a} - \mathbf{w}(\mathbf{x}_n)\|}{d(\mathbf{a}, M)} - 1 \\ &\leq F(\mathbf{x}_n) + \frac{\|\mathbf{a} - \mathbf{x}_n\|}{d(\mathbf{a}, M)} + \left(\frac{1}{d(\mathbf{a}, M)} - \frac{1}{d(\mathbf{x}_n, M)} \right) \|\mathbf{w}(\mathbf{x}_n) - \mathbf{x}_n\|. \end{aligned}$$

Using the continuity of the distance $d(\cdot, M)$ from Lemma 13.2, we obtain $F(\mathbf{a}) \leq \liminf F(\mathbf{x}_n)$.

In the same way we have:

$$F(\mathbf{x}_n) \leq \gamma^{-1}(\mathbf{w}(\mathbf{a})) + \frac{\|\mathbf{x}_n - \mathbf{w}(\mathbf{a})\|}{d(\mathbf{x}_n, M)} - 1 \leq F(\mathbf{a}) + \frac{\|\mathbf{x}_n - \mathbf{a}\|}{d(\mathbf{x}_n, M)} + \left(\frac{1}{d(\mathbf{x}_n, M)} - \frac{1}{d(\mathbf{a}, M)} \right) \|\mathbf{w}(\mathbf{a}) - \mathbf{a}\|$$

hence $\limsup F(\mathbf{x}_n) \leq F(\mathbf{a})$. Thus F is continuous and the proof is over. \square

The following lemma has a quite obvious 'proof by drawing', but it's rigorous argument is based on Brouwer's fixed point theorem.

Lemma 13.4. *Let K be the rectangle $\{[x, y] : a \leq x \leq b, c \leq y \leq d\} \subset \mathbb{R}^2$. Assume that we have two arcs $\gamma, \phi : [-1, 1] \mapsto K$, $j \in \{1, 2\}$, such that $\gamma(-1)$ belongs to the left side $\{[a, y] : y \in [c, d]\}$ of K , $\gamma(1)$ belongs to the right side $\{[b, y] : y \in [c, d]\}$ of K , $\phi(-1)$ belongs to the upper side $\{[x, d] : x \in [a, b]\}$ of K , and $\phi(1)$ belongs to the lower side $\{[x, c] : x \in [a, b]\}$ of K . Then the two arcs must cross each other, i.e. there exist $s, t \in [-1, 1]$ such that $\gamma(t) = \phi(s)$.*

Proof. Denote by $\gamma(t) = [x(t), y(t)]$ and by $\phi(s) = [u(s), w(s)]$. Assume that the two arcs never cross. It means that the quantity:

$$N(t, s) := \max\{|x(t) - u(s)|, |y(t) - w(s)|\}, \quad [t, s] \in [-1, 1] \times [-1, 1]$$

is strictly positive. By the triangle inequality:

$$|x(t) - u(s)| - |x(t_0) - u(s_0)| \leq |x(t) - x(t_0) - u(s) + u(s_0)| \leq |x(t) - x(t_0)| + |u(s) - u(s_0)|,$$

or

$$|x(t) - u(s)| \leq |x(t_0) - u(s_0)| + \epsilon \leq N(t_0, s_0) + \epsilon$$

if $[t, s]$ is close enough to $[t_0, s_0]$, due to the continuity of x and u . In a similar way we can prove $|y(t) - w(s)| \leq N(t_0, s_0) + \epsilon$, thus by taking the maximum we obtain $N(t, s) \leq N(t_0, s_0) + \epsilon$. By symmetry we must also have the inequality $N(t_0, s_0) \leq N(t, s) + \epsilon$ hence N is continuous.

Since N is continuous, positive and defined on a compact set, it must have a positive minimum. Thus $1/N(t, s)$ is also continuous on $[-1, 1] \times [-1, 1]$. Define:

$$f : [-1, 1] \times [-1, 1] \mapsto [-1, 1] \times [-1, 1], \quad f(t, s) := \left[-\frac{x(t) - u(s)}{N(t, s)}, -\frac{y(t) - w(s)}{N(t, s)} \right].$$

Due to our assumptions, $x(-1) = a$, $w(-1) = d$, $x(1) = b$ and $w(1) = c$. The function f is continuous and defined on a convex body. According to Brouwer's fixed point theorem, it must

have a fixed point. Note that the range of f belongs in fact to the boundary of the square. Thus if $f([t_0, s_0]) = [t_0, s_0]$ is a fixed point of f , we must either have $|t_0| = 1$ or $|s_0| = 1$.

If $t_0 = 1$, then we would have $1 = -\frac{x(1)-u(s_0)}{N(1,s_0)} = -\frac{b-u(s_0)}{N(1,s_0)} \leq 0$, impossible. If $t_0 = -1$ we would have $-1 = -\frac{x(-1)-u(s_0)}{N(-1,s_0)} = -\frac{a-u(s_0)}{N(-1,s_0)} \geq 0$, again impossible.

If $s_0 = 1$ we would have $1 = -\frac{y(t_0)-w(1)}{N(t_0,1)} = -\frac{y(t_0)-c}{N(t_0,1)} \leq 0$, impossible. If $s_0 = -1$ we would have $-1 = -\frac{y(t_0)-w(-1)}{N(t_0,-1)} = -\frac{y(t_0)-d}{N(t_0,-1)} \geq 0$, again impossible.

Thus f cannot have fixed points, which shows that our assumption on the positivity of N was false. □

The next lemma says that if an arc starts inside a closed rectangle and ends outside it, then it must cross the boundary.

Lemma 13.5. *Let K be the rectangle $\{[x, y] : a \leq x \leq b \text{ and } c \leq y \leq d\} \subset \mathbb{R}^2$. Assume that $\gamma : [0, 1] \mapsto \mathbb{R}^2$ is an arc such that $\gamma(0) \in \text{Int}(K)$ and $\gamma(1) \notin K$. Then there exists $0 < c < 1$ such that $\gamma(c) \in \partial K$ and $\gamma(t) \in \text{Int}(K)$ for all $0 \leq t < c$.*

Proof. We start by noting that K is closed, the interior of K is given by

$$\text{Int}(K) = \{[x, y] : a < x < b \text{ and } c < y < d\},$$

the exterior of K is

$$K^c = \{[x, y] : x < a \text{ or } b < x \text{ or } y < c \text{ or } d < y\},$$

and the boundary is $\partial K = K \setminus \text{Int}(K)$.

Denote by A the subset of the interval $[0, 1] \subset \mathbb{R}$ defined by:

$$A := \{0 \leq t \leq 1 : \gamma(s) \in \text{Int}(K), \forall 0 \leq s \leq t\}.$$

In words, if $t \in A$, then all the points of the arc γ corresponding to previous parameter values $s \leq t$ lie in the open rectangle. Since A is bounded, it has a supremum which we denote by c . Denote by $[x(t), y(t)] := \gamma(t)$. Since $a < x(0) < b$ and $c < y(0) < d$, and because x and y are continuous functions, the previous strict inequalities will remain true in a neighborhood of 0. This shows that $c > 0$. Moreover, since c is the supremum of A , there exists a sequence $\{t_n\}_{n \geq 1} \subset A$ such that $\lim_{n \rightarrow \infty} t_n = c$. This means in particular that $a < x(t_n) < b$ and $c < y(t_n) < d$ for all n , and by taking the limit using the continuity of x and y at c we obtain $a \leq x(c) \leq b$ and $c \leq y(c) \leq d$. In other words, $\gamma(s) \in \text{Int}(K)$ if $0 \leq s < c$ and $\gamma(c) \in K$.

Assume without loss of generality that $\gamma(1) = [\xi_1, \xi_2] \in K^c$ with $\xi_1 < a$. Since $a \leq x(c)$, we have that $c < 1$. Moreover, there exists N large enough such that $c + 1/n < 1$ for all $n \geq N$. Because $c + 1/n$ is not an element of A , we may find some $0 \leq s_n \leq c + 1/n$ such that $\gamma(s_n) := [u_n, w_n] \notin \text{Int}(K)$. Moreover, we must have $c \leq s_n$ because we know that for all $t < c$ we have $\gamma(t) \in \text{Int}(K)$. Thus $c \leq s_n \leq c + 1/n$. In other words, $s_n \rightarrow c$ and $\gamma(s_n) = [u_n, w_n] \notin \text{Int}(K)$. Hence at least one of the following four possibilities must occur: $u_n \leq a$ or $u_n \geq b$ or $w_n \leq c$ or $w_n \geq d$. There must exist a subsequence $\{s_{n_k}\}_{k \geq 1}$ such that exactly one of the above four inequalities is satisfied for all $k \geq 1$. Without loss of generality, assume that $u_{n_k} \leq a$ for all $k \geq 1$. Since $s_{n_k} \rightarrow c$, due to the continuity of γ at c we must have that the first component of $\gamma(c)$ must obey the same inequality as the first component of $\gamma(s_{n_k})$. Thus $\gamma(c) \notin \text{Int}(K)$. But we proved before that $\gamma(c) \in K$. Thus $\gamma(c) \in K \setminus \text{Int}(K) = \partial K$ and we are done. □

13.2 The main theorem

If $\phi : S^1 \mapsto \mathbb{R}^2$ is a homeomorphism which maps the unit circle into the plane, then the image $J := \phi(S^1)$ is called a Jordan curve. In other words, a Jordan curve is a simple closed path which can be parametrized by

$$[0, 2\pi] \ni t \mapsto \phi(\cos(t), \sin(t)) \in \mathbb{R}^2.$$

A Jordan curve is bounded and closed, thus compact.

Theorem 13.6. (*Jordan curve theorem*). *Let J be any Jordan curve. Then the set $\mathbb{R}^2 \setminus J$ is open in \mathbb{R}^2 , has exactly two connected components (one bounded and the other one unbounded), and J is their boundary.*

Proof. We start by proving that $\mathbb{R}^2 \setminus J$ is not connected, i.e. it has at least two connected components. Assume that $\mathbb{R}^2 \setminus J$ is connected. According to Lemma 13.1, since $\mathbb{R}^2 \setminus J$ is open (because J is closed), it must be path connected. The strategy is to construct an 'inner' point \mathbf{x}_i which cannot be joined with the points situated outside some large ball which contains J .

Let us construct this point \mathbf{x}_i . The map

$$\mathbb{R}^4 \supset J \times J \ni [\mathbf{x}, \mathbf{y}] \mapsto \|\mathbf{x} - \mathbf{y}\| \in \mathbb{R}$$

is continuous and defined on a compact set. Thus there exist \mathbf{x}_l and \mathbf{x}_r in J which maximize this distance, i.e. $\|\mathbf{x} - \mathbf{y}\| \leq \|\mathbf{x}_r - \mathbf{x}_l\|$ for all $\mathbf{x}, \mathbf{y} \in J$. Denote by $\overline{\mathbf{x}_r \mathbf{x}_l}$ the straight segment joining the two points, and consider the two infinite lines L_l and L_r which are perpendicular on $\overline{\mathbf{x}_r \mathbf{x}_l}$ and pass through \mathbf{x}_l and \mathbf{x}_r respectively. No point of L_l other than \mathbf{x}_l , and no point of L_r other than \mathbf{x}_r can belong to J , otherwise $\|\mathbf{x}_r - \mathbf{x}_l\|$ would not be maximal. Thus J belongs to the strip generated by the two lines. Moreover, because J is bounded, we can build a closed rectangle K which includes J in its interior and has two parallel sides included in L_l and L_r .

Without loss of generality, we may assume that $\mathbf{x}_l = [-1, 0]$, $\mathbf{x}_r = [1, 0]$ and $K = \{[x, y] : -1 \leq x \leq 1, -10 \leq y \leq 10\}$. The curve J has exactly two points in common with K , and they are \mathbf{x}_r and \mathbf{x}_l . These two points split J into two arcs: J_u and J_d . Without loss of generality we may assume that J_u starts at \mathbf{x}_r and ends at \mathbf{x}_l with the trigonometric orientation, while J_d starts at \mathbf{x}_l and ends at \mathbf{x}_r with the same trigonometric orientation.

The segment linking the top point $T = [0, 10]$ with the bottom point $B = [0, -10]$ is denoted by \overline{TB} . We note that J_u and \overline{TB} are two arcs which must cross at least in one point, due to Lemma 13.4. Denote by $\mathbf{y}_{max,u}$ the point of $J_u \cap \overline{TB}$ with the largest second coordinate, i.e. the crossing point closest to T . Denote by $\mathbf{y}_{min,u}$ the point of $J_u \cap \overline{TB}$ with the lowest second coordinate, i.e. the crossing point closest to B . Note that it can happen that $\mathbf{y}_{max,u} = \mathbf{y}_{min,u}$.

In the same way, the arc J_d and the segment \overline{TB} must cross. Denote by $\mathbf{y}_{max,d}$ the point of $J_d \cap \overline{TB}$ with the largest second coordinate, i.e. the crossing point closest to T . Denote by $\mathbf{y}_{min,d}$ the point of $J_d \cap \overline{TB}$ with the smallest second coordinate, i.e. the crossing point closest to B .

Define the 'inner' point which we talked about to be $\mathbf{x}_i := (\mathbf{y}_{min,u} + \mathbf{y}_{max,d})/2$. Clearly, \mathbf{x}_i is not an element of J and belongs to $\mathbb{R}^2 \setminus J$. If $\mathbb{R}^2 \setminus J$ were connected, we can join \mathbf{x}_i with any other point from outside K , since $K^c \subset \mathbb{R}^2 \setminus J$. According to Lemma 13.5, such an arc must cross the boundary of the rectangle K in some point \mathbf{w} . This \mathbf{w} can be neither \mathbf{x}_r nor \mathbf{x}_l , since they belong to J .

If the second coordinate of \mathbf{w} is negative, then consider the arc starting at T , continued with a straight segment to $\mathbf{y}_{max,u}$, continued with the part of J_u between $\mathbf{y}_{max,u}$ and $\mathbf{y}_{min,u}$, then by the straight segment to \mathbf{x}_i , then by the arc linking \mathbf{x}_i with \mathbf{w} , and then we continue on the boundary of K until we reach B . In this way we constructed an arc in K starting at T and ending at B which has no common points with J_d , contradicting Lemma 13.4.

If the second coordinate of \mathbf{w} is positive, then consider the arc starting at B , continued with a straight segment to $\mathbf{y}_{min,d}$, then continued with the part of J_d between $\mathbf{y}_{min,d}$ and $\mathbf{y}_{max,d}$, then with a straight segment to \mathbf{x}_i , then with the arc from \mathbf{x}_i to \mathbf{w} , and then on the boundary of K until we reach T . In this way we constructed an arc in K linking B with T which does not cross J_u , again a contradiction. Thus $\mathbb{R}^2 \setminus J$ is not connected.

Up to now we know that there exists exactly one unbounded connected component (which contains K^c), and at least one bounded 'inner' component. The next result is about the boundary of each such connected component: it says that if U is a connected component of $\mathbb{R}^2 \setminus J$, then U is open in \mathbb{R}^2 and the boundary $\partial U = \overline{U} \setminus U$ equals J .

The first observation is that $\partial U \subset J$; if this were not true, then there exists some point \mathbf{x} in \overline{U} which belongs neither to U nor to J , hence it must be an element of some other connected component W . But then \mathbf{x} is an inner point of W and must be isolated from U , contradiction with our assumption that \mathbf{x} belongs to the closure of U .

It could happen though that ∂U is strictly included in but not equal with J . In this case, there exists an arc $M \subset J$ such that $\partial U \subset M$. We will show that this leads to a contradiction.

We first assume that U is a bounded connected component. Consider a closed ball $D := \overline{B_R(\mathbf{x}_o)}$ where \mathbf{x}_o is some inner point of U , and $R > 0$ is sufficiently large such that the circle ∂D belongs to K^c , thus outside U . Clearly, $M \subset \overline{U} \subset D$. According to Lemma 13.3, there exists a continuous map $g : D \mapsto M$ such that $g(\mathbf{x}) = \mathbf{x}$ on M . Define the map $q : D \mapsto D \setminus \{\mathbf{x}_o\}$ given by $q(\mathbf{x}) = g(\mathbf{x}) \in M$ if $\mathbf{x} \in \overline{U}$ and $q(\mathbf{x}) = \mathbf{x}$ if $\mathbf{x} \in D \setminus U$. Note that q is well defined because the 'dangerous' points of $\overline{U} \cap (D \setminus U)$ are included in M where $g(\mathbf{x}) = \mathbf{x}$. Moreover, q is continuous, and \mathbf{x}_o is never in its range. Let $t : D \setminus \{\mathbf{x}_o\} \mapsto \partial D$ be the natural retraction, i.e. the map which sends $\mathbf{x} \in D \setminus \{\mathbf{x}_o\}$ into the point on ∂D obtained from the intersection of ∂D with the ray starting from \mathbf{x}_o and going through \mathbf{x} . Let $a : \partial D \mapsto \partial D$ be the antipodal map, i.e. the map which sends a point of ∂D into the diametrically opposed point. Now define the map $r : D \mapsto D$ given by $r = a \circ t \circ q$. We note that r is continuous, and its range is ∂D . Brouwer's fixed point theorem says that r must have a fixed point, which can only be on the boundary ∂D . But $q(\mathbf{x}) = \mathbf{x}$ if $\mathbf{x} \in \partial D \subset U^c$, and $t(\mathbf{x}) = \mathbf{x}$ on the boundary. The antipodal map a prohibits the existence of a fixed point on the boundary for r , which leads to a contradiction. Thus $\partial U = J$ if U is bounded.

If W is the unbounded connected component and M is an arc containing the boundary of W , then we can use the point \mathbf{x}_o and the disk D previously considered in order to define $q : D \mapsto D \setminus \{\mathbf{x}_o\}$ by $q(\mathbf{x}) = \mathbf{x}$ if $\mathbf{x} \in \overline{W}$ and $q(\mathbf{x}) = g(\mathbf{x}) \in M$ if $\mathbf{x} \in D \setminus W$. From here the rest of the argument is identical, and leads to a contradiction. Thus $\partial W = J$.

Let us recapitulate what we know until now: there is exactly one unbounded component, at least one bounded, and every connected component has its boundary equal with J . The last thing we have to prove is that there are no other bounded components besides the component U_i which contains the point \mathbf{x}_i defined when we proved that $\mathbb{R}^2 \setminus J$ is not connected.

Now let us assume that there exists a bounded component $W \neq U_i$. The points \mathbf{x}_r and \mathbf{x}_l belong to the boundary of W , hence both of them are limits of sequences of points from W . In particular, there exists $\tilde{\mathbf{x}}_r \in W$ such that its first component is larger than $1/2$, and there exists $\tilde{\mathbf{x}}_l \in W$ such that its first component is smaller than $-1/2$. Because W is path connected, there exists a path $\widehat{\tilde{\mathbf{x}}_l \tilde{\mathbf{x}}_r} \subset W$.

Now consider the path \widehat{TB} starting from T , continued with the straight segment joining T with $\mathbf{y}_{max,u}$, continued with the arc of J between $\mathbf{y}_{max,u}$ and $\mathbf{y}_{min,u}$, then with the straight segment (containing \mathbf{x}_i) between $\mathbf{y}_{min,u}$ and $\mathbf{y}_{max,d}$, continued with the arc of J between $\mathbf{y}_{max,d}$ and $\mathbf{y}_{min,d}$, and finally continued with the straight segment between $\mathbf{y}_{min,d}$ and B . We see that all the points of \widehat{TB} belong either to J , to U_i or to the unbounded connected component. It means that \widehat{TB} and $\widehat{\tilde{\mathbf{x}}_l \tilde{\mathbf{x}}_r} \subset W$ cannot have common points, and this contradicts Lemma 13.4. □