

# Complex Analysis Notes for MP3

Horia Cornean, d.6/12/2010.

## 1 Some typical exam exercises

**Exercise 1.1.** Find all complex solutions to the equation  $e^{z^2} = 1$ .

**Solution.** We know that the exponential function is  $2\pi i$  periodic, thus  $z^2$  must be of the form  $2\pi iN$  with  $N \in \mathbb{Z}$ . There are three possibilities for  $N$ :

1. If  $N = 0$ , then the only solution is  $z = 0$ ;
2. For each  $N > 0$ , let us solve the equation  $z^2 = 2\pi iN = 2\pi N e^{i\pi/2}$ . For each positive  $N$  we find two solutions:

$$z_k = (2\pi N)^{1/2} e^{i(\pi/4 + \pi k)} = (-1)^k (2\pi N)^{1/2} e^{i\pi/4} = (-1)^k (\pi N)^{1/2} (1 + i), \quad k \in \{0, 1\}.$$

3. For each  $N < 0$ , let us solve the equation  $z^2 = -2\pi i|N| = 2\pi|N|e^{-i\pi/2}$ . This gives other two solutions:

$$z_k = (2\pi|N|)^{1/2} e^{i(-\pi/4 + \pi k)} = (-1)^k (2\pi|N|)^{1/2} e^{-i\pi/4} = (-1)^k (\pi N)^{1/2} (1 - i), \quad k \in \{0, 1, 2\}.$$

**Exercise 1.2.** Let  $f(z) = |z|^2 + \bar{z}$ , where  $z = x + iy$ .

1. Find two real functions  $u$  and  $v$  such that  $f(z) = u(x, y) + iv(x, y)$  for all  $z$ .
2. Is  $f$  analytic?

**Solution.**

1. We have  $\bar{z} = x - iy$  and  $|z|^2 = x^2 + y^2$ , thus  $u(x, y) = x + x^2 + y^2$  and  $v(x, y) = -y$ .
2. The function is not analytic, because the Cauchy-Riemann equations are not satisfied. For example,  $\partial_x u = 1 + 2x$  is not identically equal with  $\partial_y v = -1$ .

**Exercise 1.3.** Let  $f(z) = \bar{z}$ , where  $z = x + iy$ . Let  $\gamma$  be a circle of radius 1, centred at  $z_0 = 1 + i$ , and oriented anti-clockwise. Show that the path integral

$$\int_{\gamma} f(z) dz = 2\pi i.$$

Is this result in contradiction with Cauchy's integral theorem?

**Solution.** We can parameterize the circle as  $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$ ,  $\gamma(t) = z_0 + e^{it}$ . Here  $\gamma'(t) = ie^{it}$  and  $z_0 = \sqrt{2}e^{i\pi/4}$ . Then we have

$$\begin{aligned} \int_{\gamma} f(z) dz &= \int_0^{2\pi} f(\gamma(t)) \gamma'(t) dt = i \int_0^{2\pi} \overline{\sqrt{2}e^{i\pi/4} + e^{it}} e^{it} dt \\ &= i \int_0^{2\pi} (\sqrt{2}e^{-i\pi/4} e^{it} + 1) dt = 2\pi i. \end{aligned} \tag{1.1}$$

Thus the integral of  $f$  on a closed path is not zero. This is possible because  $f$  is not analytic, thus Cauchy's theorem is not contradicted.

**Exercise 1.4.** Find the convergence radius of the power series  $\sum_{n \geq 0} (1+i)^n \frac{2n^3+i}{(n+i)^2} z^n$ .

**Solution.**

Let  $a_n = (1+i)^n \frac{2n^3+i}{(n+i)^2}$ . This implies that  $a_{n+1} = (1+i)^{n+1} \frac{2(n+1)^3+i}{(n+1+i)^2}$ . Then we have:

$$\frac{a_n}{a_{n+1}} = \frac{1}{1+i} \frac{(2n^3+i)(n+1+i)^2}{(2(n+1)^3+i)(n+i)^2} = \frac{1}{1+i} \frac{(2+i/n^3)(1+(1+i)/n)^2}{(2(1+1/n)^3+i/n^3)(1+i/n)^2}, \quad n \geq 1.$$

Thus the radius of convergence is:

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \frac{1}{\sqrt{2}}.$$