


8. Session: Vectors

Def.: A vector in the plane (2D-vector) is an ordered pair of real numbers

$$\vec{v} = (a, b).$$

The numbers a and b are called coordinates or components of \vec{v} . A scalar is a real number.


Geometric interpretation



Representatives for \vec{v} .


Def. The length of $\vec{v} = (a, b)$ is

$$\|\vec{v}\| = \sqrt{a^2 + b^2}.$$



Ex. $\vec{v} = (-1, 2)$

$$\|\vec{v}\| = \sqrt{(-1)^2 + 2^2} = \sqrt{5}$$




Let $\vec{u} = (u_1, u_2)$, $\vec{v} = (v_1, v_2)$ and let c be a scalar.

Def. $\vec{u} = \vec{v}$ if and only if $u_1 = v_1$ and $u_2 = v_2$.




Def. $\vec{u} + \vec{v} = (u_1 + v_1, u_2 + v_2)$

• $c\vec{u} = (cu_1, cu_2)$

Geometric interpretation



or



Theorem: $\|c\vec{u}\| = |c| \|\vec{u}\|$.

Proof: $\|c\vec{u}\| = \|(cu_1, cu_2)\| = \sqrt{(cu_1)^2 + (cu_2)^2} = \sqrt{c^2u_1^2 + c^2u_2^2}$
 $= \sqrt{c^2(u_1^2 + u_2^2)} = \sqrt{c^2} \cdot \sqrt{u_1^2 + u_2^2} = |c| \|\vec{u}\|$ q.e.d.

Def.: The zero vector

The negative of \vec{u}

Subtraction

Division by scalar $c \neq 0$

$$\vec{0} = (0, 0)$$

$$-\vec{u} = (-1)\vec{u} = (-u_1, -u_2)$$


$$\vec{u} - \vec{v} = \vec{u} + (-\vec{v}) = (u_1 - v_1, u_2 - v_2)$$

$$\frac{\vec{u}}{c} = \frac{1}{c}\vec{u} = \left(\frac{u_1}{c}, \frac{u_2}{c}\right)$$

Remark: Two points $P = (p_1, p_2)$ and $Q = (q_1, q_2)$ determine a vector \vec{QP} . One can compute its coordinates as follows:

$$\vec{QP} = P - Q = (p_1 - q_1, p_2 - q_2)$$

That is, $\vec{QP} = \vec{OP} - \vec{OQ}$:



The distance between P and Q is

$$|QP| = \|\vec{QP}\| = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2}$$

Ex. For $P = (3, 2)$ and $Q = (1, -1)$, we have

$$\vec{QP} = P - Q = (3, 2) - (1, -1) = (3 - 1, 2 - (-1)) = (2, 3).$$

Theorem: For vectors $\vec{a}, \vec{b}, \vec{c}$ and scalars r, s one has

$$(1) \vec{a} + \vec{b} = \vec{b} + \vec{a}$$

$$(4) (r+s)\vec{a} = r\vec{a} + s\vec{a}$$

$$(2) \vec{a} + (\vec{b} + \vec{c}) = (\vec{a} + \vec{b}) + \vec{c}$$

$$(5) (rs)\vec{a} = r(s\vec{a})$$


$$(3) r(\vec{a} + \vec{b}) = r\vec{a} + r\vec{b}$$

(Proof: write out in coordinates.)

Def. \vec{u} is called a unit vector if $\|\vec{u}\| = 1$.

Theorem (Normalization)

If $\vec{w} \neq \vec{0}$, then $\frac{\vec{w}}{\|\vec{w}\|}$ is a unit vector in the same direction as \vec{w} .



Proof: $\frac{1}{\|\vec{w}\|} > 0$ and $\left\| \frac{\vec{w}}{\|\vec{w}\|} \right\| = \left\| \frac{1}{\|\vec{w}\|} \vec{w} \right\| = \left| \frac{1}{\|\vec{w}\|} \right| \|\vec{w}\| = \frac{\|\vec{w}\|}{\|\vec{w}\|} = 1$.
q.e.d.


Def. The standard basis vectors are

$$\vec{i} = (1, 0) \quad \text{and} \quad \vec{j} = (0, 1).$$

Remark:


$$\vec{u} = (u_1, u_2) = u_1 \vec{i} + u_2 \vec{j}$$

Proof: $u_1 \vec{i} + u_2 \vec{j} = u_1(1, 0) + u_2(0, 1) = (u_1, 0) + (0, u_2) = (u_1, u_2) = \vec{u}$.
q.e.d.




$$(2, 3) = 2\vec{i} + 3\vec{j}$$

3D - space



Right-handed
coordinate system




Coordinates of a
point

Def. A vector in 3D-space is an ordered triple of real numbers

$$\vec{v} = (a, b, c).$$

Geometric interpretation



Let $\vec{u} = (u_1, u_2, u_3)$, $\vec{v} = (v_1, v_2, v_3)$ be vectors and let c be a scalar.

$$\|\vec{u}\| = \sqrt{u_1^2 + u_2^2 + u_3^2} \quad \text{length}$$

$$\vec{u} + \vec{v} = (u_1 + v_1, u_2 + v_2, u_3 + v_3)$$

$$c\vec{u} = (cu_1, cu_2, cu_3)$$

As for vectors in the plane, one has

- $\|c\vec{u}\| = |c| \|\vec{u}\|$

- Two points $P = (p_1, p_2, p_3)$ and $Q = (q_1, q_2, q_3)$ determines a vector \vec{QP} .




$$\vec{QP} = P - Q.$$

$$|QP| = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2 + (p_3 - q_3)^2}.$$

Standard basis vectors

$$\vec{i} = (1, 0, 0), \quad \vec{j} = (0, 1, 0), \quad \vec{k} = (0, 0, 1).$$



$$\vec{u} = (u_1, u_2, u_3) = u_1 \vec{i} + u_2 \vec{j} + u_3 \vec{k}.$$